

Report of the Federal Task Force on
High Risk Pregnancies and
Prenatal Record Systems



Canada

REPORT OF THE
FEDERAL TASK FORCE ON HIGH RISK PREGNANCIES
AND PRENATAL RECORD SYSTEMS

Également disponible en français sous
le titre "Rapport du groupe fédéral de
travail sur les grossesses à risque
élevé et les systèmes d'enregistrement
des données prénatales"

SUMMARY OF RECOMMENDATIONS OF THE
FEDERAL TASK FORCE ON HIGH RISK PREGNANCY
AND PRENATAL RECORD SYSTEMS

Foreword

This report is not a statement of federal government policy but represents an information base for further considered courses of action and potential policy development. It is hoped that this report will prove to be a useful guide for those professionals and para-professionals involved in providing pre-natal and peri-natal care to expectant mothers.

SUMMARY OF RECOMMENDATIONS OF THE FEDERAL TASK FORCE ON HIGH
RISK PREGNANCY AND PRENATAL RECORD SYSTEMS

The following are the recommendations of the Federal Task Force:

RECOMMENDATIONS (SECTION II)

1. Each province in Canada should be encouraged to develop a prenatal record form for use by physicians undertaking obstetrical care.
2. The prenatal forms devised in each province should as far as possible be uniform in design and concept and should include a means of risk evaluation of the expectant mother using accepted risk predictors detailed in the body of this report.
3. Provision should be made for inclusion of the original prenatal record form in the mother's hospital record, a copy to be included in the neonatal hospital record and a copy to be retained in the physician's office record.
4. Provincial efforts to develop regionalized programs of perinatal care which have resulted in triage of high risk pregnancies according to their care requirements are laudable and should be encouraged throughout Canada.
5. Although the primary intent of the prenatal record form is to facilitate care of the pregnant woman by her physician, consideration should be given during the course of form design to inclusion of certain basic items of information which, if uniformly entered and collected, could be used for data retrieval and analysis.

RECOMMENDATIONS (SECTION III)

6. Properly designed studies into the validity of risk scoring systems should be encouraged and supported.

7. Risk assessment of the pregnant woman is of proven value and should be universally applied.
8. Properly conducted and controlled studies must be instituted to demonstrate the effect of length of work and type of occupation on pregnancy.
9. The cost implications of alternative work proposals to pregnant women and society must be evaluated.
10. Studies should be undertaken on the effect of toxic substances in the workplace, and their effects on the pregnant working woman and her fetus.
11. Prospective studies should be conducted to clarify the relationship of early pregnancy bleeding to pregnancy outcome, with special emphasis on congenital anomalies.
12. Well-controlled, prospective studies of the effect of nutritional supplementation and diet on the outcome of pregnancy are necessary.
13. A strong research effort should be made to elucidate the pathophysiology of pregnancy induced hypertension and its relationship to poor maternal outcome.
14. Adequately controlled prospective studies into post-term pregnancies must be carried out, with particular emphasis on fetal and maternal morbidity.
15. The Task Force recommends the continued support of prenatal care programs, with special emphasis on the detection of pregnancies at risk.
16. Continuing emphasis should be given to the humanization of the birthing process to maintain and improve psychological satisfaction of this family experience. However, the fundamental guiding principle of a responsible patient care provider must always be to maintain and protect the physical well-being of the mother and her infant.

17. Information is urgently required to assess the potential for long-term effects of ultrasound on the fetus.
18. Until such studies of long-term effects of ultrasound are completed, the use of routine ultrasound without specific indications in pregnancy should be discouraged.
19. Educational standards for ultrasonographers are required.
20. Properly designed prospective studies of antepartum and intrapartum fetal heart monitoring should be implemented to establish the validity of specific applications.
21. Properly designed trials should be carried out to delineate side effects on mother and fetus and attempt made to validate whether there is improvement in outcome by use of tocolytic agents.
22. Maternal-fetal transport should be undertaken whenever possible in preference to transport of the distressed neonate.
23. Programs of maternal transport within the framework of a regionalized system should be implemented uniformly throughout Canada in preference to neonatal transport.
24. Because of potential hazardous long-term effects, glucocorticoids to prevent respiratory distress in the neonate should only be administered when it is felt that their definite short-term benefits are essential.
25. Whereas the role of the nurse in reproductive care is recognized in many countries, it has not yet been agreed upon in Canada, and therefore the Committee proposes the establishment of a National Task Force to address this important issue. The "core" members of the Task Force should include clinical perinatal nurses plus obstetrical and pediatric perinatologists. Additional representatives should be appointed from appropriate medical and nursing professional bodies.

26. Neonatal morbidity should be precisely defined and its appropriateness to reflect the quality and effects of perinatal care delineated and evaluated.

RECOMMENDATION (SECTION IV)

27. The Task Force recommends that the foregoing definitions of uniform terminology for maternal and perinatal statistics be adopted for use throughout Canada. It is further recommended that these definitions be forwarded to the appropriate federal and provincial authorities for further discussion with a view to their implementation.

RECOMMENDATION (SECTION V)

28. An ad hoc committee of experts on prenatal and perinatal care be established and be charged with follow-up of the Report of the Canadian Commission for the International Year of the Child and of this Task Force report for the specific purpose of ensuring that Canadians attain optimal outcome of pregnancy as previously defined.

TABLE OF CONTENTS

	<u>Page</u>
SUMMARY OF RECOMMENDATIONS OF THE FEDERAL TASK FORCE ON HIGH RISK PREGNANCY AND PRENATAL RECORD SYSTEMS	
SECTION I. GENERAL INTRODUCTION.....	1
A. BACKGROUND.....	1
B. PARTICIPANTS.....	2
C. PREAMBLE.....	3
D. OBJECTIVES OF THE TASK FORCE.....	4
E. METHOD OF STUDY.....	4
SECTION II. PRENATAL RECORDS.....	6
A. CONTENT.....	6
B. UNIFORMITY.....	6
C. INCLUSION IN HOSPITAL RECORDS.....	8
D. PERINATAL CARE PROGRAMS.....	9
E. DEFINITIONS OF RISK PREGNANCY.....	10
a) Risk Markers.....	12
b) Outcome Markers.....	13
c) Intervention Markers.....	13
SECTION III. EVALUATION OF PREDICTORS, INTERVENTIONS AND OUTCOME VARIABLES.....	15
INTRODUCTION.....	15
A. EPIDEMIOLOGIC METHODOLOGY.....	15
B. ACCEPTED RISK PREDICTORS.....	17
1) Previous Obstetrical Performance.....	17
2) Maternal Medical Factors.....	17
3) Socioeconomic Factors.....	17
4) Fetal and Maternal Factors Arising During Pregnancy.....	18
5) Labour and Delivery Factors.....	18
C. CONTROVERSIAL PREDICTORS.....	19
1) Risk Scoring.....	19
2) Length of Work and Occupation During Pregnancy.....	20
3) Toxic Exposures and Congenital Abnormalities.....	21
4) Early Antepartum Bleeding and Pregnancy Outcome.....	22
5) Nutritional Supplement.....	22

	<u>Page</u>
6) Blood Pressure Elevation: Maternal Significance.....	23
7) Post-Term Pregnancy.....	23
D. CONTROVERSIAL INTERVENTIONS.....	23
1) Prenatal Care; Providers and Programs.....	23
2) Childbirth Education Programs.....	24
3) Ultrasound.....	25
4) Intrapartum Electronic Heart Rate Monitoring.....	26
5) Tocolytics.....	27
6) Maternal versus Neonatal Transport.....	27
7) Amniotic Fluid Assays.....	28
8) Glucocorticoids and Enhanced Fetal Lung Maturity.....	29
9) Intrapartum Care - Obstetrician, General Practitioner or Nurse in Reproductive Care.....	30
10) Mortality and Morbidity.....	31
SECTION IV. REPORT OF THE SUBCOMMITTEE ON UNIFORM MATERNAL AND PERINATAL TERMINOLOGY.....	32
INTRODUCTION.....	32
RECOMMENDED UNIFORM DEFINITIONS.....	32
SECTION V. FUTURE OF THE TASK FORCE ON HIGH RISK PREGNANCY.....	38
APPENDIX 1. REFERENCES.....	39
APPENDIX 2. TASK FORCE MEMBERS AND CONSULTANTS.....	73
APPENDIX 3. REPORT OF SUBCOMMITTEE ON THE PRENATAL RECORD.....	78
APPENDIX 4. LITERATURE REVIEW: FETAL HEART RATE MONITORING, DR. RUTH MILNER.....	90
APPENDIX 5. PROVINCIAL PRENATAL RECORD FORMS.....	110

I. GENERAL INTRODUCTION

A. BACKGROUND

There have been many significant advances in reproductive care in recent years. Application of new knowledge and techniques has resulted in progressive reduction in perinatal mortality rates in Canada to values which are now comparable with the best in the world. It is obvious however that mortality rates should not be the only criterion upon which the success of a nation's program for perinatal health care is judged: morbidity rates and client satisfaction are two other important criteria which must be taken into account if optimal outcomes are to be attained in what is a physiological and joyous event for the majority of the population.

This Federal Task Force on High Risk Pregnancies and Prenatal Record Systems was established in February 1981 by the Health Services Directorate of the Health Services and Promotion Branch. Its formation arose through a series of steps initiated by the Canadian Institute of Child Health (C.I.C.H.) in 1978 to attempt to improve the outcome of perinatal care in Canada. In that year studies were initiated, one result of which was formation of the Canadian Coalition for the Prevention of Handicaps Through Improved Prenatal and Perinatal Care. Other outcomes included preparation of a position paper and a book entitled "Priority of Priorities"⁽¹⁾ copies of which were sent to all Provincial ministers and their deputies and to the Federal Minister of Health and Welfare.

In 1979 the Report of the Canadian Commission for the International Year of the Child recommended that governments "adopt a policy making maternal and child health a priority for the next five years"⁽²⁾.

In 1980, Monique Bégin, Minister of Health and Welfare, in a letter to the Chairman of the Board of the C.I.C.H., William Cochrane, M.D., F.R.C.P.(C), commending the Institute for its "role in stressing the need for improved perinatal care", suggested further discussions with her department to plan and develop a coordinated program.

These discussions prompted the formation of this Task Force in 1981. Its terms of reference, listed by M.M. Law, M.D., F.R.C.P.(C), Assistant Deputy Minister, Health Services and Promotion Branch, Health and Welfare, Canada, were as follows:

1. To discuss, develop and process a set of guidelines which will identify various maternal conditions which will aid health professionals in recognizing cases of high risk pregnancy.
2. To develop a prenatal record system which would have the potential for universal use and applicability.

B. PARTICIPANTS

The following were members of the Task Force:

Dr. Jack H. Walters, Chairman	Dr. Thomas Martin
Dr. Graham Chance	Dr. John Milligan
Dr. James Dunne	Dr. Ruth Milner
Dr. Martine Gagnon	Dr. Ian Morrison
Dr. Emily Hamilton	Dr. Leo Peddle
Dr. Karen Kaufman	Mrs. Shirley Post
Mrs. Joyce (MacDonald) Kaye	Dr. Donald Rae
Dr. Bernard Leduc	Dr. D. Smith
Dr. Robert H. Lennox	Dr. H. Stewart
Dr. T. MacLachlan	

Consultants in Epidemiology and a clinical consultant group who provided valuable comments are listed in Appendix 2. This report was prepared by the members of the Committee who are practising health professionals and scientists with particular expertise and interest in perinatal care. The report is submitted to government but was not prepared by government. Valuable support from Dr. Lennox and his secretarial staff was provided to arrange the meetings, transcribe and translate the minutes and numerous task force documents, and in the preparation and printing of this final report. The document is intended for the use of practising health professionals who provide care for pregnant women and their infants. It is also planned to distribute it to Health Services planners at all levels of government, who require guidelines and standardized definitions for the development of improved systems of perinatal health care.

C. PREAMBLE

The concept of perinatal care has evolved from a time when there were no uniformly accepted definitions or specialized units, through to the development of an internationally accepted concept of high risk pregnancy and Perinatal Care units for high risk pregnancy patients and their infants, made available through regionalized programs.

The literature review and those citations in the bibliography in this report are limited to references concerning high risk pregnancies which have been published over the past 10 years. They are but a sample of the many publications in this important field, and include 368 references which have been reviewed by Dr. Ruth Milner and her group.

D. OBJECTIVES OF THE TASK FORCE

The specific objectives of the Task Force were projected into the above historical sequences. A number of the provinces have already developed systems whose stated objectives are to reduce perinatal mortality and morbidity. These provinces have developed record systems, interventions as well as personnel and facilities to accomplish these objectives. The Task Force studied the possibility of developing a set of uniform definitions for Canada, and the development of a set of "core markers" which could be considered for use by each province in modifying their existing prenatal record systems, if indicated or needed.

The conclusions and recommendations of the Task Force which follow are not intended to coerce or mandate the development of a single prenatal record form, method of management or system of health care of prenatal patients in any province. They are intended to establish standards for possible comparison of similar conditions, methods of management and outcomes in the perinatal field. Only through the cooperation of the members of the professional associations and the federal and provincial health care systems, will improvements evolve. As a group of concerned professionals, we urge that perinatal care be given the proper priority at all levels over the next years. We trust that this report will aid in such an evolution.

E. METHOD OF STUDY

The three main tasks of the Task Force for the study were allocated to three separate subcommittees which were asked to review the literature and prepare reports which could be studied and approved by the main Task Force.

The Subcommittee on Prenatal Records was chaired by Dr. I. Morrison. The report of this Committee is summarized in Section II and given in detail in Appendix 3.

The second Subcommittee under the chairmanship of Dr. B. Leduc and Dr. J. Milligan was given the more difficult task of defining high risk pregnancies. After their initial meeting they met with the whole committee, and the following motion was adopted in recognition of our concern over certain essential aspects of modern care.

"Recognizing the concern of the Health Care Professions and of the general public over the apparently high incidence of 'High Risk Pregnancies', it is moved that the validity of currently used predictors of unfavourable outcome of viable pregnancies and the basis upon which the outcomes are identified, be evaluated."

Dr. Ruth Milner agreed to serve on the Task Force to provide the expertise and supervision for these evaluations.

The Subcommittee on Definitions under the chairmanship of Dr. Leo Peddle, developed the section which is included in the report entitled "Uniform Maternal and Perinatal Terminology". This section has been forwarded to the recently-formed National Committee on Perinatal Terminology which has representatives from Statistics Canada as well as this Task Force to be used as the basis for a set of uniform definitions which could be acceptable to provincial registrars and federal authorities.

II. PRENATAL RECORDS

A. CONTENT

In the medical or social history of a pregnant mother and findings on physical and laboratory examination, there may be "markers" or "factors" which indicate potential problems for the mother or child. These "markers" may be identified at an early stage of pregnancy or may develop at any time throughout the pregnancy, up to the time of onset of labour and delivery.

The factors sampled, and assessed by family physicians, obstetricians and nurses in the management of the pregnant mother undergoing prenatal care, may be recorded on specifically designed forms called prenatal record forms. These forms are used by the health care team for documenting ongoing care of pregnant mothers. They are usually designed so that they are easily identified as separate and specific records for use in offices, clinics and hospital medical record departments.

The history of the development of the prenatal record and the increasing sophistication of its content over the past half century, mirrors the development of the specialty of perinatology. There was a reduction of international maternal and perinatal mortality rates as techniques to detect and manage major obstetrical and neonatal complications became known. To a limited extent this reduction was accompanied by a lowering of morbidity in these same risk situations. However, as outcomes have improved perinatal mortality has approached its "irreducible minimum" and reduced morbidity of perinatal origin must nowadays be regarded as the prime focus of effective care.

B. UNIFORMITY

To improve the identification of pregnant women "at risk" and to provide a standard guideline for the health care team

documenting the numerous factors evaluated throughout the prenatal period for all pregnant mothers, groups of health professionals developed comprehensive prenatal records. Initially these records were utilized only for ambulatory patients in offices and clinics, but gradually hospitals, cities, regions and eventually provinces developed uniform prenatal records. During their evolution, prenatal records have undergone many user trials. In general the following changes can be identified:

1. The format of the record became more standardized.
2. Certain items of information were cited in specific areas of the record (e.g. obstetrical history or last menstrual period).
3. Sections of the record were developed for notations of follow-up or management (e.g. weight, blood pressure, fetal health).
4. The items or factors recognized to be required for prenatal management became accepted (e.g. blood group, Rhesus type and antibodies, Rubella titre).
5. New research findings used in this sorting or identification of risk became incorporated (e.g. symphysis-fundus height measurements, ultrasound).

Specific areas for recording of cervical cytology and for listing of pertinent personal, medical, family or social factors became established as integral aspects of the prenatal record. Concurrent with development of prenatal records, the principle that certain pregnancies benefit from special care has become more widely established and accepted.

C. INCLUSION IN HOSPITAL RECORDS

With improvements in prenatal data acquisition and recognition of the significance of prenatal events for postnatal outcomes, there was a recognition of the importance of the prenatal record in the management of the mother and her infant, to act as a vehicle to transmit information to the labour, delivery and postnatal wards of the hospital in which she was to give birth.

At first only the blood group, Rhesus type and serology report were seen to be required. Eventually blood pressure, total weight gain, hemoglobin, investigations and treatments carried out during the prenatal period and other information was realized to be needed by the perinatal team. As a result, the record is now usually in multicopy format. In most instances, one copy is sent to the hospital and the original is kept in the doctor's office or clinic. In some provinces an additional copy is sent to the neonatal chart. In many hospitals in Canada, the by laws of the hospital oblige all practitioners to forward to the hospital a copy of the prenatal record for each patient who is to be admitted in labour to that hospital. Most provinces now have standardized prenatal records for this purpose.

Recommendation #1. Each province in Canada should be encouraged to develop a prenatal record form for use by physicians undertaking obstetrical care.

Recommendation #2. The prenatal forms devised in each province should as far as possible be uniform in design and concept and should include a means of risk evaluation of the expectant mother, using accepted risk predictors detailed in the body of this report.

Recommendation #3. Provision should be made for inclusion of the original prenatal record form in the mother's hospital record, a copy to be included in the neonatal hospital record and a copy to be retained in the physician's office record.

A more detailed discussion of the content, design and purpose of the prenatal record is found in Appendix 3.

D. PERINATAL CARE PROGRAMS

The dramatic reduction of maternal and perinatal deaths and the concomitant reduction in morbidity for both mothers and their newborns over the past 15 years in Canada is not only due to improved housing, food or transportation. It is also due to the recognition that a proportion of all pregnant women, probably less than 25 per cent, have increased risk perhaps due to some unrecognized problem(s), that require(s) these women to be managed differently from the remainder. Over the past 15 years, these "risk" patients have been identified, more appropriately managed, and their perinatal death and morbidity rates reduced as a result of application of various interventions by health care teams.

A major component of this improvement has been accomplished through education of mothers-to-be to the recognition that their pregnancy is indeed "at risk" and that there are possibilities for improving the outcome for many of them.

The progressive improvement in perinatal statistics in all regions of Canada attests to the impact of these programs.

The system of health care delivery has been modified over the same period of time so as to ensure availability of special care for those mothers with particular needs. Facilities have been developed on a regional basis to provide teams of personnel specifically trained to care for the identified high risk patient, her fetus and newborn. These teams have been based in specially designed and equipped perinatal units which are the designated centres for regional educational programs and for referral of women with complicated pregnancies. In conjunction with the other institutions and professionals in the region, the facilities serve as resource centres to provide the expertise necessary to bring about improved care. This phenomenon of "regionalization of medical care" is well established for other special medical problems, including

cancer, trauma, renal dialysis and genetics. Such an approach has been shown to ensure equitable availability of costly programs and to improve outcome for these special medical conditions.

Recommendation #4. Provincial efforts to develop regionalized programs of perinatal care which have resulted in triage of high risk pregnancies according to their care requirements are laudable and should be encouraged throughout Canada.

For the small number of instances in which either mother or baby dies or in which there is a continuing disability of the mother or child, anguish and suffering may be prolonged. Moreover, the health care cost for continuing care of damaged infants and children is a financial burden for the family and for society for many years. Further reduction of perinatal mortality and morbidity among the offspring of those pregnant mothers who are identified at risk, will thus be to the benefit of society. Achievement of this aim will require better understanding of causal factors, many of which originate prenatally. Record linkage of information concerning the social, economic and educational background of the woman and her partner is important in long-term assessment of risk factors.

Recommendation #5. Although the primary intent of the prenatal record form is to facilitate care of the pregnant woman by her physician, consideration should be given during the course of form design to inclusion of certain basic items of information which, if uniformly entered and collected, could be used for data retrieval and analysis.

E. DEFINITIONS OF RISK PREGNANCY

A consensus was reached by the Task Force that the definition of high risk pregnancy was dependent upon the sum total of the markers which individually are associated with increased risk during a pregnancy. There were three groups of markers identified by the Task Force as follows:

a) Risk Markers

These are factors which have been accepted and are currently being used in clinical practice as indicators of some degree of increased risk in any pregnancy. An example is maternal diabetes.

b) Outcome Markers

Unfavourable outcome events which can be identified and are valid as the current markers for at least this pregnancy. Examples would be maternal sepsis or intra-ventricular hemorrhage.

c) Intervention Markers

These are techniques or factors which have a positive or negative influence on the management and/or outcome of the pregnancy. An example would be the measurement of lecithin-sphingomyelin ratio in the amniotic fluid as a predictor of pulmonary maturity.

Based on a consensus of clinical experience, the following factors thought to influence the pregnant patient, the fetus or the neonate, were listed by the Committee and placed in the following categories, according to priority, for methodological evaluation of their literature:

1. Those factors which were so well recognized or "obvious" that no further research or statistical assessment was felt necessary to validate them as markers (low priority for study).
2. Those factors with high priority to establish validity as markers of high risk pregnancy.

The listing by category which included Risk, Outcome and Intervention markers is as follows:

a) Risk Markers

1 (Obvious)

- 1) low birth weight infant
- 2) smoking
- 3) previous fetal or neonatal death
- 4) drugs (non-medical)
- 5) medical disorders
- 6) antibodies (Rh, Rubella, etc.)
- 7) alcohol ingestion
- 8) gestational diabetes
- 9) no prenatal care
- 10) multiple pregnancy
- 11) premature rupture of membranes
- 12) prematurity
- 13) abnormal position of fetus
- 14) abnormal presentation of fetus
- 15) anesthetic coverage and resuscitation
- 16) scalp sampling of fetal blood

2 (Validity to be Established)

- 1) time of first attendance for prenatal care
- 2) urinary tract infection
- 3) therapeutic abortion history
- 4) pre-pregnancy weight
- 5) pre-pregnancy height
- 6) socioeconomic status
- 7) nutrition
- 8) alpha-feto protein
- 9) age - under 20 years
- over 35 years
- 10) occupation - mother (not specifically father)
- 11) validity of standard form used
- 12) validity of risk screening form used
- 13) bleeding
- 14) hypertension (two readings 20 minutes apart)
- 15) pregnancy longer than 42 weeks
- 16) labelling of patient high risk
- 17) screening with non-stress test
- 18) prenatal classes attendance
- 19) place of birth (home vs. hospital)
- 20) size of hospital (either in or out of functioning regional hospital system)

b) Outcome Markers

1 (Obvious)

- 1) respiratory distress syndrome
- 2) intraventricular hemorrhage
- 3) sepsis
- 4) genetic and developmental abnormalities
- 5) broncho-pulmonary dysplasia

2 (Validity to be Established)

- 1) anemia
- 2) bilirubin
- 3) fractures
- 4) small for gestational age
- 5) seizures
- 6) cerebral palsy
- 7) mental retardation
- 8) neurological abnormalities at discharge
- 9) low Apgar (< 7 at 5 minutes)
- 10) metabolic acidosis (cord pH < 7.2)

c) Intervention Markers

1 (Obvious)

- 1) skilled neonatal transport
- 2) maternal-fetal transport

2 (Validity to be Established)

- 1) who does the delivery (G.P., Obs., midwife)
- 2) maternal-fetal transport versus neonatal transport
- 3) bed rest
- 4) operative delivery (Cesarean, forceps)
- 5) induction
- 6) glucocorticoids
- 7) tocolytic agents

Funding was made available to Dr. Milner and her staff for a research project to review the above list of markers which were discussed and categorized by the members of the Task Force. The objective was to make a critical assessment of the validity of the scientific methodology used in evaluation in published studies relating to these markers and to explore areas where their use was less than optimal.

The section on methodology outlines the stringent review methods used by Dr. Milner and her group. The section entitled Intervention and Outcome Variables outlines the results of these extensive reviews. The comments and recommendations of the Task Force for areas of further study and development are based on these studies.

III. EVALUATION OF PREDICTORS, INTERVENTIONS AND OUTCOME VARIABLES

INTRODUCTION

Pregnancy is a natural process, however complications may develop and thus incur risks to the mother or child. The natural reproductive process ensures survival of the species, but on an individual basis does not guarantee a problem-free pregnancy nor a healthy newborn. The intent of modern perinatal care is to optimize the birth of an infant with full potential while maintaining minimal risk to the mother.

In order to be effective in preventing problems, risk detection must be prospective. The establishment of predictive factors has been dependent largely upon retrospective evaluation. Some complications in pregnancy may be immediate and unpredictable, for example the occurrence of premature rupture of the membranes at 27 weeks, or cord prolapse in an otherwise apparently healthy pregnancy. In contrast, past events, such as the history of premature rupture of membranes at 27 weeks in a previous pregnancy, indicate a potential for risk in a current pregnancy, yet in the majority of patients a healthy outcome will ensue.

This Task Force has examined factors believed to be associated with increased risk of adverse outcome which may be used to alert the physician to actual or potential problems. Some, which were considered non-controversial and generally accepted, are only listed (Section B). Other factors claimed in the literature to be predictive and/or to be appropriate interventions, but which are controversial, have been extensively scrutinized with regard to their proven epidemiologic validity as predictors of problems or interventions considered beneficial to patient management (Sections C and D).

A. EPIDEMIOLOGIC METHODOLOGY

The criteria for assessment of the quality of the evidence were based on those used by the Canadian Task Force on the Periodic Health Examination⁽³⁾.

Three aspects were reviewed. They were:

- the current burden of suffering;
- the validity and acceptability of the manoeuvre used to identify the risk of condition;
- the effectiveness of the resulting intervention.

Only the latter two aspects were reviewed in detail.

To obtain the appropriate literature, a Medline Search for the previous two years was run for each risk marker or intervention, and recent articles were solicited from colleagues. From articles cited, other references were culled from the bibliographies attached to each and a library was thus assembled around every marker. A positive decision was made not to review literature prior to 1975 because it was felt that recent evidence would be more valuable. Only in rare cases was the search taken further; either because the best articles were published earlier and still reflected current evidence, or because there was a dearth of recent articles.

To assess the quality of evidence contained in the articles, the criteria for judging diagnostic tests or evaluating the effectiveness of interventions were taken from the series of critical appraisal articles which appeared in the Canadian Medical Association Journal in 1981 under the title of Clinical Epidemiology Rounds⁽⁴⁾. An attempt was made to cover the three aspects already mentioned as fully as possible but lack of time, data and appropriate references precluded a complete review, especially in regard to the current burden of suffering.

An example of the application of the literature appraisal methods used is given in the review of the literature on Intrapartum Fetal Monitoring in Appendix 4.

B. ACCEPTED RISK PREDICTORS

1. Previous Obstetrical Performance

Past obstetrical history has a strong predictive value for the current pregnancy. This contribution is unfortunately not available for nulliparas and even in multiparas is not absolutely predictive.

Low birth weight infants (<2.5 kg)

Perinatal death

Habitual abortions (3 or more consecutive)

Birth injury

Congenital anomalies

Genetic disease

Grand multiparity

Previous Cesarean section

2. Maternal Medical Factors

Recognition of these factors is important because appropriate intervention will minimize complications.

Medical disorders in pregnancy (e.g. diabetes, hypertension, etc.)

Isoimmunization

Family history of genetic disease

Maternal age (≤ 16 years, or ≥ 35 years)

Rubella susceptibility

Uterine surgery or abnormality

3. Socioeconomic Factors

Extremes of unfavourable socioeconomic and cultural factors produce detrimental effects on the pregnancy.

Poverty and starvation
No prenatal care
Alcoholism
Heavy smoking
Drug abuse

4. Fetal and Maternal Factors Arising During Pregnancy

These factors affirm the importance of early and ongoing prenatal care for the detection, assessment and management of actual or potential problems in pregnancy.

Multiple pregnancy
Intrauterine growth retardation
Concurrent infection (e.g. Rubella, herpes, toxoplasmosis, syphilis)
Fetal disease (e.g. hydrops, genetic diseases, congenital anomalies)
Polyhydramnios
Oligohydramnios
Maternal disease (e.g. pregnancy induced hypertension (PIH), gestational diabetes, anemia)
Antepartum bleeding
Isoimmunization

5. Labour and Delivery Factors

Previously identified risk factors may well persist at the time of onset of labour; however, additional and non-predicted complications may arise during labour and delivery. These may be associated with major problems and can require urgent and immediate attention.

Preterm labour
Premature rupture of membranes
Prolonged labour
Malpresentation and/or Malposition
Prolapsed cord
Traumatic birth
Passage of Meconium
Intrapartum asphyxia
Intrapartum bleeding
Neonatal asphyxia
Congenital anomalies interfering with neonatal adaptation
(e.g. diaphragmatic hernia)

C. CONTROVERSIAL PREDICTORS

1. Risk Scoring is the numerical weighting of individual and combined risk factors which are added up to a cumulative score.

Risk Assessment is the clinical evaluation of the significance of individual risk factors.

Risk scoring has been reported to correlate with perinatal outcome. However, when subjected to strict epidemiological analysis, the validity of these correlations is questionable.

Properly designed studies have not yet been published which tested the hypothesis that risk scoring by itself modified perinatal outcome. However the proper application of clinical assessment of risk factors (Section B, 1-5) has been associated with improved perinatal outcome for individuals.

The educational value, especially to the health care provider, in the use of risk scoring and/or risk assessment, when applied to a population of pregnant women, is recognized (5-33).

COMMENTS ON RECOMMENDATIONS

It is apparent that although there are many factors which may potentially increase the risk of adverse outcome of pregnancy, there is urgent need for further research to determine whether these factors represent real or imagined risk, or whether certain interventions have real or imagined benefit. Evaluation of the significance of some factors may be precluded because studies designed to test them may be too complex and impractical or may contain major ethical obstacles. Likewise, certain accepted intervention procedures of imprecisely defined value also may be difficult to withdraw.

Recommendation #6. Properly designed studies into the validity of risk scoring systems should be encouraged and supported.

Recommendation #7. Risk assessment of the pregnant woman is of proven value and should be universally applied.

2. Length of Work and Occupation During Pregnancy

Currently there are no adequate data nor published reports to answer the questions of the effect of length of work or the type of occupation on pregnancy and its outcome.

This area has become of major importance as the number of women of childbearing age entering the work force has increased consistently over the past number of years. Moreover, legislation, union contracts, and job benefits for working pregnant women make the answers to these questions of a great practical importance. The main conclusion of the American College of Obstetricians and Gynecologists in their report for the U.S. Department of Health and Education states that "The normal woman with an uncomplicated pregnancy and a normal fetus in a job that presents no greater potential hazards than those encountered in normal daily life in the community, may continue to work without interruption until the onset of labour and may resume working several weeks after an uncomplicated delivery". However, while accepting

the concept of work throughout pregnancy, this conclusion reveals the lack of factual data necessary to advise pregnant women.

The economic implications, both of working restrictions and of cost of benefits need to be delineated with respect to the individual and society. The effects of work within the home on pregnancy and its outcome, the relationship of stress, socio-economic factors, etc. to employment and pregnancy must be clarified. The relationship of non-employment related activities such as crafts, athletics, etc. have not been evaluated and need to be outlined.

Recent trends in Canada concerning the employment of pregnant women, and the financial and social implications of such changes are a major issue in our society today (34-41).

Recommendation #8. Properly conducted and controlled studies must be instituted to demonstrate the effect of length of work and type of occupation on pregnancy.

Recommendation #9. The cost implications of alternative work proposals to pregnant women and society must be evaluated.

3. Toxic Exposures and Congenital Abnormalities

Precise identification of potential responsible agents, exposure levels, and susceptible periods during pregnancy are lacking in most studies. Spontaneous abortion is reported as increased in operating room personnel, in steel smelting workers, and in those working in styrene or rayon production. Congenital abnormalities are reported in higher prevalence among children of gardeners, switchboard operators, reinforced plastic production, steel smelting workers, pharmaceutical laboratory workers, operating room workers and workers using organic solvents. Intensive studies in these women and in those working in other fields such as those working with Visual Display Terminals are needed to clarify exposure risks in pregnancy.

Recommendation #10. Studies should be undertaken on the effect of toxic substances in the workplace, and their effects on the pregnant working woman and her fetus.

4. Early Antepartum Bleeding and Pregnancy Outcome

Early antepartum bleeding prior to 20 weeks does warn of some increased risk of late pregnancy complications, namely, premature birth, congenital anomalies, and increased perinatal mortality. The Ontario Perinatal Mortality Study ⁽⁶⁴⁾ showed a premature birth rate of 5.9% in patients with no early bleeding and 14.5% in patients with early bleeding. The perinatal mortality rate was also increased from 22.9 to 65.2 in those without bleeding compared to those with bleeding. The only studies showing an increased congenital anomaly rate were retrospective and therefore included a strong recall bias⁽⁴²⁻⁵³⁾.

Recommendation #11. Prospective studies should be conducted to clarify the relationship of early pregnancy bleeding to pregnancy outcome, with special emphasis on congenital anomalies.

5. Nutritional Supplement

Although much is written of the importance of good nutrition in pregnancy and there is some correlation between low pre-pregnancy weight and low weight gain with low birth weight babies, nutritional supplementation alone has not been shown to have a direct effect on pregnancy outcome. Many study design problems exist in published articles, such as failure to ascertain whether the supplement was actually eaten, whether the intended supplement was actually taken in addition to usual diet or if it simply substituted for the normal diet. Such study defects generally preclude adequate evaluation of the effects of supplemental nutrition⁽⁵⁴⁻⁷²⁾.

Recommendation #12. Well controlled, prospective studies of the effect of nutritional supplementation and diet on the outcome of pregnancy are necessary.

6. Blood Pressure Elevation: Maternal Significance

Pregnancy may unmask latent hypertension but information on the early and late consequences to the mother is scant. Severe pregnancy induced hypertension remains one of the leading causes of maternal mortality⁽⁷³⁻⁹³⁾.

Recommendation #13. A strong research effort should be made to elucidate the patho-physiology of pregnancy induced hypertension and its relationship to poor maternal outcome.

7. Post-Term Pregnancy

Because perinatal mortality is relatively low after post-term pregnancy, the use of this marker for statistical analysis is limited. However, descriptive studies report increased incidence of induction of labour, meconium stained amniotic fluid, and Cesarean section, and indicate the need for further investigation with an emphasis on morbidity⁽⁹⁴⁻¹⁰²⁾.

Recommendation #14. Adequately controlled prospective studies into post-term pregnancies must be carried out with particular emphasis on fetal and maternal morbidity.

D. CONTROVERSIAL INTERVENTIONS

1. Prenatal Care: Providers and Programs

Very few studies have adequately examined the relationship between the training and qualifications of prenatal care providers and pregnancy outcome. The difficulties inherent in such studies almost preclude their satisfactory completion (e.g. the frequency of prenatal examination and the timing of the first visit are closely related to socio-economic factors which themselves affect pregnancy outcome).

It was interesting to note that in one study in family practice in which prenatal care was randomly allocated to either physician or nurse practitioner, no difference was shown in outcome.

Studies in rural communities have demonstrated significant improvement in outcome when prenatal care programs conducted by nurse-midwife were introduced where no programs previously existed. From the evidence reviewed, it is clear that the difficulties inherent in conducting studies of the effectiveness of prenatal care programs or their provision by different personnel, have almost precluded their satisfactory completion⁽¹⁰³⁻¹¹⁵⁾.

Recommendation #15. The Task Force recommends the continued support of prenatal care programs with special emphasis on the detection of pregnancies at risk.

2. Childbirth Education Programs

Recent emphasis on childbirth education programs and their possible impact on perinatal outcome has been a source of major controversy. For this reason, it was felt important to validate the suggested beneficial effects of these programs. Unfortunately, their content and evaluation criteria differ amongst all studies, making validity assessment impossible.

A second question has related to the efficacy of these particular interventions, and again the reports in the literature are fraught with difficulties because of methodological problems. However the use of analgesia, anesthesia and forceps is generally decreased in those women who have participated in such programs.

There is no difference in length of labour, frequency of complications, birth weight, or Apgar score but there does tend to be a greater psychological satisfaction with the birth experience.

Unfortunately, none of the studies examined cost benefits or effectiveness. No long-range benefits have been demonstrated. It does not appear that possible disadvantages such as unrealistic expectations or aims have actually been examined. It is difficult to assess effectiveness of the intervention as the goals addressed by the various approaches are diverse⁽¹¹⁶⁻¹³⁷⁾.

Recommendation #16. Continuing emphasis be given to the humanization of the birthing process to maintain and improve psychological satisfaction of this family experience. However, the fundamental guiding principle of a responsible patient care provider must always be to maintain and protect the physical well-being of the mother and her infant.

3. Ultrasound

Ultrasound examinations are helpful in the detection of anomalies, assessment of gestational age, multiple gestation, placental localization, and fetal heart rate and movement. The benefits of ultrasound scanning vary with the particular function expected from it. As far as fetal growth is concerned, there seems to be very good evidence that ultrasound is a useful adjunct to clinical identification and assessment of intra-uterine growth retardation.

Ultrasound is also useful to determine fetal age when it is used rationally and early in pregnancy. While there have been few prospective controlled studies, it appears clear to the Committee that ultrasound is necessary prior to amniocentesis⁽¹³⁸⁻²⁰¹⁾.

Recommendation #17. Information is urgently required to assess the potential for long-term effects of ultrasound on the fetus.

Recommendation #18. Until such studies of long-term effects of ultrasound are completed, the use of routine ultrasound without specific indications in pregnancy should be discouraged.

Recommendation #19. Educational standards for ultrasonographers are required.

4. Intrapartum Electronic Heart Rate Monitoring

Fetal Heart Rate Monitoring was introduced to help to identify the fetus in distress, so that appropriate intervention could be considered. Most studies show that electronic fetal monitoring has a high negative predictive value, that is, a normal tracing is very likely to be associated with a non-asphyxiated infant. Unfortunately, an abnormal tracing does not correlate as well with poor outcome and therefore must be interpreted with all other available data. The National Institute of Child Health and Development's Consensus Development Task Force on Antenatal Diagnosis and the Society of Obstetricians and Gynecologists of Canada suggest that the use of electronic monitors be strongly considered in "high risk" patients. There is no acceptable evidence to indicate any beneficial effects of electronic fetal monitoring in the pregnancy with no identified risk.

Because of the complexity and importance of evaluation of intrapartum fetal monitoring, the concerned reader is referred to Appendix 4.

Antepartum and intrapartum fetal heart rate monitoring are widely utilized in Canada today and are of apparent value. However, the literature is lacking in adequate studies to assess some of the applications of these techniques⁽²⁰²⁻²⁴³⁾.

Recommendation #20 Properly designed prospective studies of antepartum and intrapartum fetal heart monitoring should be implemented to establish the validity of specific applications.

5. Tocolytics

Premature delivery is the most important contributor to perinatal mortality and morbidity. Prevention and treatment of premature labour are therefore of particular importance.

Effectiveness of tocolysis is difficult to assess due to methodological problems (e.g. in defining premature labour, in obtaining placebo medication with similar side effects to minimize observer bias, sample size, standardization of terminal objectives). Examination of reports shows that the objective of treatment was not always similar (e.g. to delay delivery for 24 to 48 hours, to reduce incidence of RDS or deaths, or to increase the number of term deliveries)⁽²⁴⁴⁻²⁸³⁾.

Recommendation #21 Properly designed trials should be carried out to delineate side effects on mother and fetus and attempt to validate whether there is improvement in outcome by use of tocolytic agents.

6. Maternal versus Neonatal Transport

Precise assessment of the benefit of neonatal versus maternal transport is not possible since no randomized trial of transport versus retention of the mothers or neonates in non-specialized centres has been carried out. All reported studies have major methodological problems, not the least of which relates to the fact that in neonatal transport, the patients are selected before transport by evidence of illness, or

excluded where hopeful survival does not appear to be present. By contrast, selection of mothers for maternal transport is based on either current or potential problems. It seems apparent that fetuses requiring intensive and specialized care around birth would benefit from transfer to centres where that expertise is readily available. For this reason, although desirable, it seems that a randomized trial would be unethical.

Maternal transport for the sake of fetal or maternal potential problems has been accepted by the profession and all provincial health ministries in Canada. The concept of transporting the fetus at risk in utero to the perinatal team at an appropriate centre has reached a consensus in theory. This Committee strongly supports this concept.

Implementation of the concept of maternal-fetal transport is still inadequate because of application problems and lack of support at various levels of responsibility⁽²⁸⁴⁻³⁰²⁾.

Recommendation #22. Maternal-fetal transport should be undertaken whenever possible in preference to transport of the distressed neonate.

Recommendation #23. Programs of maternal transport within the framework of a regionalized system should be implemented uniformly throughout Canada in preference to neonatal transport.

7. Amniotic Fluid Assays

Measurement of amniotic fluid optical density is of proven value in the management of isimmunization as is the use of amniotic fluid for assessment of risk of certain genetic abnormalities in appropriately selected mothers. A large number of studies over the last several years on many of the

constituents of amniotic fluid have not proved to be of clinical use relative to fetal maturity. The major reviews at the present time relate to measurement of amniotic fluid phospholipids and especially the relationship of the Lecithin-Sphingomyelin (L/S) ratio to prediction of fetal lung maturity. There is a large body of evidence which supports the validity of these tests in relation to outcome. A review of the literature strongly supported this conclusion. These reviews did not include the disorders where the L/S ratio is of lessened value, for example in the diabetic pregnancy.

There was no review of the usefulness of the more recently recognized constituents of amniotic fluid such as phosphatidylglycerol (PG). However the Committee recognizes that this is an area of great promise⁽³⁰³⁻³¹¹⁾.

8. Glucocorticoids and Enhanced Fetal Lung Maturity

The Committee wished to make a statement concerning this controversial therapy in the prevention of respiratory distress syndrome (RDS) in the premature infant, but time constraints precluded detailed literature review.

Two randomized clinical trials showed that glucocorticoids reduced the incidence of respiratory distress syndrome (RDS) but there has not been a clear demonstration of decreased perinatal mortality. There is no evidence of significant adverse short-term effects but long-term side effects potentially exist and are being evaluated (reference N.I.H. study).

Recommendation # 24. Because of potential hazardous long-term effects, glucocorticoids to prevent respiratory distress in the neonate should only be administered when it is felt that their definite short-term benefits are essential.

9. Intrapartum Care - Obstetrician, General Practitioner or Nurse in Reproductive Care

There are no Canadian studies available as the practice of Obstetrics by other than licensed physicians is not legal in Canada. Studies from the United States suggest that nurse-midwife practices are a safe, economical and acceptable means of providing obstetrical care to selected groups of women. The evidence is inconclusive to the extent that findings are confounded by:

1. A variety of practice settings from home to free-standing and hospital-based birth centres and regular hospital delivery suites.
2. Differing eligibility criteria for the nurse of reproductive care clients (and other selection factors).
3. Joint practice management and transfer to physician care at various points in the pregnancy.

The outcome of the studies can be classified under three broad areas, namely Safety, Acceptability and Cost. None of the studies demonstrated increased perinatal mortality as a result of supervision by a nurse in reproductive care. No study specifically addressed acceptability although all studies concluded that on the basis of continued demand and referral the system was acceptable. Despite initial reservations on the part of some physicians, all studies report eventual acceptance of the nurse in reproductive care service.

There are no Canadian data which examine the role of the family doctor vis-à-vis the obstetrician. Specialists are functioning as primary care physicians for many pregnant women and many family doctors do not practise obstetrics.

It would seem that there is a complementary role for each of the three: nurse in reproductive care, family physician and obstetrician. Any system will be most efficient when personnel are utilized appropriately according to their level of training⁽³¹²⁻³²²⁾.

Recommendation #25. Whereas the role of the nurse in reproductive care is recognized in many countries, it has not yet been agreed upon in Canada, and therefore the Committee proposes the establishment of a National Task Force to address this important issue. The "core" members of the Task Force should include clinical perinatal nurses plus obstetrical and pediatric perinatologists. Additional representatives should be appointed from appropriate medical and nursing professional bodies.

10. Mortality and Morbidity

There is general recognition that perinatal mortality may not be an adequate indicator to reflect quality of perinatal care or to highlight areas where preventive activity is required. However, neonatal morbidity has not been precisely defined. A study of some area of morbidity (e.g. neonatal seizures, mental retardation and cerebral palsy) might in part accomplish such a goal.

Recommendation # 26. Neonatal morbidity should be precisely defined and its appropriateness to reflect the quality and effects of perinatal care delineated and evaluated.

IV. REPORT OF THE SUBCOMMITTEE ON UNIFORM MATERNAL AND PERINATAL TERMINOLOGY

INTRODUCTION

At an early meeting of the Task Force on High Risk Pregnancy it was requested that uniform definitions of perinatal terminology be developed. Considerable work has been carried out in this area during the past 10 years. The Subcommittee used the following documents for developing recommendations for standardization of maternal and perinatal terminology:

1. Recommendations of the Perinatal Medicine Committee of the Society of Obstetricians and Gynecologists of Canada and the Fetus and Newborn Committee of the Canadian Pediatric Society, October 1974.
2. Information paper entitled "Hospital Perinatal Committee" published by the Canadian Institute of Child Health in 1981 and printed in "Dimensions in Health Care Services", 1979.
3. Williams Obstetrics, Sixteenth Edition, by Pritchard and MacDonald.

RECOMMENDED UNIFORM DEFINITIONS

Birth

Birth is the complete expulsion or extraction from its mother of a fetus weighing 500 g or more, irrespective of the gestational age.

Livebirth

Livebirth is the birth of a fetus, irrespective of gestational age, weighing 500 g or more which after its complete expulsion or extraction,

breathes or shows any other evidence of life, such as beating of the heart, pulsation of the umbilical cord, or definite movement of the voluntary muscles.

Stillbirth (Fetal Death)

Stillbirth is the birth of a fetus weighing 500 g or more, irrespective of gestational age, who shows no sign of life after birth.

Abortion

Abortion is the complete expulsion or extraction from its mother of a fetus or embryo weighing less than 500 g irrespective of gestational age, whether there is evidence of life or not, and whether the abortion was spontaneous or induced.

Preterm Infant

An infant born before the 37th completed week of gestation (less than 259 days).

Term Infant

An infant born during a period beginning at the 37th completed week of gestation and extending to the 42nd completed week (259 to 293 completed days).

Post Term Infant

An infant born after 42 completed weeks of gestation (294 days or more).

Low Birthweight

A fetus or infant weighing less than 2500 g but 500 g or more at birth.

Full Birthweight (full size)

A fetus or infant weighing 2500 g or more at birth.

Birthweight

Refers to the first weight of the fetus or newborn obtained after birth. This weight should be measured within the first hour of life before

significant postnatal weight loss has occurred.

Recommended birthweight groupings:

500 g, up to and including 749 g
750 g, up to and including 999 g
1000 g, up to and including 1249 g
1250 g, up to and including 1499 g
1500 g, up to and including 1999 g
2000 g, up to and including 2499 g
2500 g, up to and including 2999 g
3000 g, up to and including 3499 g
3500 g, up to and including 3999 g
4000 g, up to and including 4499 g
4500 g, up to and including 4999 g
5000 g, and over

Early Neonatal Period

The interval after birth but during the first 7 completed days of life.

Early Neonatal Death

Death of a liveborn infant weighing 500 g or more at birth, during the first 7 completed days of life.

Early Neonatal Mortality Rate

The early neonatal mortality rate is the number of early neonatal deaths of infants weighing 500 g or more, per 1000 liveborn infants.

Late Neonatal Period

The interval beginning after 7 and up to 28 completed days of life.

Late Neonatal Death

Death of a liveborn infant weighing 500 g or more at birth after 7 but before 28 completed days of life.

Late Neonatal Mortality Rate

The late neonatal mortality rate is the number of late neonatal deaths of infants weighing 500 g or more, per 1000 liveborn infants.

Fetal Death (Stillbirth)

Stillbirth is the birth of a fetus weighing 500 g or more, irrespective of gestational age, showing no sign of life after birth.

Fetal Death Rate (Stillbirth Death Rate)

The stillborn infant rate is the number of stillborn infants weighing 500 g or more, per 1000 total births.

Perinatal Period

The interval from the time a fetus attains a weight of 500 g through to the first 7 completed days of intrauterine life.

Perinatal Death

Stillborn and liveborn infants, both weighing 500 g or more who died before birth, at birth, or during the first 7 completed days of life.

Perinatal Mortality Rate (500 grams or more)

$$\frac{\text{Fetal Deaths + First Week Neonatal Deaths} \\ \text{(both 500 grams or more)}}{\text{Livebirths + Fetal Deaths} \\ \text{(both 500 grams or more)}} \times 1000$$

Standard Perinatal Period

The interval from the time a fetus attains a weight of 1000 g through the first 7 completed days of life.

Standard Perinatal Death

Stillborn and liveborn infants, both weighing 1000 g or more who died before birth, at birth, or during the first 7 completed days of life.

Standard Perinatal Mortality Rate (1000 grams or more) (international)

$$\frac{\text{Fetal Deaths} + \text{First Week Death} \\ (\text{both 1000 grams or more})}{\text{Livebirths} + \text{Fetal Deaths} \\ (\text{both 1000 grams or more})} \times 1000$$

Note: "First Week" to be interpreted as seven periods of 24 hours.

Maternal Death

Death that occurs as a result of the reproductive process and which is made up of direct maternal deaths and indirect maternal deaths.

Direct Maternal Death

Death of the mother resulting from obstetric complications in the pregnancy state, labour or puerperium and from interventions, omissions, incorrect treatment or a chain of events resulting from any of the above. (Example, exsanguination from rupture of the uterus.)

Indirect Maternal Death

Death of the mother not directly related to obstetric causes but resulting from previously existing disease, or a disease that developed during pregnancy, labour or puerperium, but which was aggravated by the maternal physiological adaptation to pregnancy. (Example, mitral stenosis.)

Non-Obstetrical Death

Death of the mother resulting from accidental or incidental causes in no way related to pregnancy. (Example, death from an airplane crash.)

Maternal Mortality Rate

The number of maternal deaths that occur as a result of the reproductive process per 100 000 live births.

It should be noted that the above definitions of maternal mortality adapted from Williams Obstetrics do not specify a gestational time period. It is suggested that for use in this definition the time from conception to 42 days postpartum be accepted.

Recommendation #27. The Task Force recommends that the foregoing definitions of uniform terminology for maternal and perinatal statistics

be adopted for use throughout Canada. It is further recommended that these definitions be forwarded to the appropriate federal and provincial authorities for further discussion with a view to their implementation.

V. FUTURE OF THE TASK FORCE ON HIGH RISK PREGNANCY

As discussed in the general Introduction, the Report of the Canadian Commission for the International Year of the Child recommended that governments "adopt a policy making maternal and child health a priority for the next five years". The Task Force on High Risk Pregnancy has welcomed this opportunity to have input into aspects of management of the prenatal period, attention to which will result in improved outcome for mothers and babies. Knowledge and attitudes to perinatal care are changing rapidly and the Task Force considers it important that the momentum for improved care attained through its establishment not be lost. The Task Force therefore recommends that an ad hoc committee on prenatal and perinatal care be established in association with the committee set up to follow-up on the recommendations of the Canadian Commission for the International Year of the Child.

Recommendation #28. An ad hoc committee of experts on prenatal and perinatal care be established and be charged with follow-up of the Report of the Canadian Commission for the International Year of the Child and this Task Force Report for the specific purpose of ensuring that Canadians attain optimal outcome of pregnancy as previously defined.

APPENDIX 1

REFERENCES

APPENDIX 1

REFERENCES

1. Canadian Institute of Child Health: "The Priority of Priorities and the Prevention of Handicap of Prenatal-Perinatal Origin", 1980.
2. Report of Canadian Commission for the International Year of the Child: "For Canada's Children - National Agenda for Action", 1979.
3. Task Force Report, Canadian Task Force on the Periodic Health Examination, CMAJ, Vol. 121, November 3, 1979.
4. Department of Clinical Epidemiology and Biostatistics, McMaster University, Health Sciences Centre: "Clinical Epidemiology Rounds", CMAJ, Vol. 124(6):703-710; 124(7):869-872; 124(8):985-990; 124(9):1156-1162, 1981.
5. Wilson, R.W. and Schiffrin, B.S.: "Is any Pregnancy Low Risk?", Obstet. Gynecol., 55(5):653, 1980.
6. Lesinski, J.: "High Risk Pregnancy", Obstet. Gynecol., 46(5):599, 1975.
7. Newcombe, R.G. and Chalmers, I.: "Assessing the Risk of Pre-Term Labour", Butterworths International Medical Reviews. Eds. Elder, M.G. and Hendricks, C.H.
8. Task Force Report, Canadian Task Force on the Periodic Health Examination, CMAJ, Vol. 121, November 3, 1979.
9. Goodwin, J.W., Dunne, J.T. and Thomas, B.W.: "Antepartum Identification of the Fetus at Risk", CMAJ, Vol. 101:58, October 1969.

10. Yeh, S.Y., Forsythe, A., Lowensohn, R.I. and Hon, E.H.: "A Study of the Relationship Between Goodwin's High Risk Score and Fetal Outcome", Am. J. Obstet. Gynecol., 127(1):50, 1977.
11. Coopland, A.T., Peddle, L.J., Baskett, T.F., Rollwagen, R., Simpson, A., and Parker, E.: "A Simplified Antepartum High Risk Pregnancy Scoring Form: Statistical Analysis of 5459 Cases", CMAJ, 116:999, May 1977.
12. Morrison, I., and Olsen, J.: "Perinatal Mortality and Antepartum Risk Scoring", Obstet. Gynecol., 53(3):362, 1979.
13. Nesbitt, R.E.L. and Aubry, R.H.: "High Risk Obstetrics", Am. J. Obstet. Gynecol., 103(7):1972, 1969.
14. Aubry, R.H. and Pennington, J.C.: "Identification and Evaluation of High Risk Pregnancy: The Perinatal Concept", Dept. of Obs/Gyn, New York, New York.
15. Papiernik, E. and Kaminski, M.: "Multifactorial Study of the Risk of Prematurity at 32 Weeks of Gestation", J. Perinat. Med., 2:30, 1974.
16. Rantakallio, P.: "Groups at Risk in Low Birth Weight Infants and Perinatal Mortality", from: Department of Pediatrics, University of Oulu, Finland, Oulu 1969.
17. Donahue, C.L. and Wan, T.T.H.: "Measuring Obstetric Risks of Prematurity: A Preliminary Analysis of Neonatal Death", Am. J. Obstet. Gynecol., 116(7):911, 1973.
18. Hobel, C.J., Hyvarinen, M.A., Okada, D.M., Oh, W.: "Prenatal and Intrapartum High Risk Screening", Am. J. Obstet. Gynecol., 117(1):1, 1973.
19. Hobel, C.J.: "Risk Assessment in Perinatal Medicine", Clinical Obstet. Gynecol., 21(2):287, 1978.

20. Winters, S., Itzkowitz, S. and Johnson, K.: "Prenatal Risk Assessment: An Evaluation of the Hobel Record in a Mount Sinai Clinic Population", The Mount Sinai J. of Med., 46(4):424, 1979.
21. Sokol, R.J., Rosen, M.G., Stojkov, J., and Chik, L.: "Clinical Application of High Risk Scoring on an Obstetric Service", Am. J. Obstet. Gynecol., 128(6):652, 1977.
22. Effer, S.B.: "Biochemical and Biophysical Indices of Fetal Risk", Clinics in Perinatology, 1(1):161, 1974.
23. Mercier, G. and Desjardins, P.D.: "Evaluation Numérique Du Risque Pendant La Grossesse", TOME 102, January 1973.
24. Rumeau-Rouquette, C., Kaminski, M., and Goujard, J.: "Prediction of Perinatal Mortality in Early Pregnancy", J. Perinat. Med., 2:196, 1974.
25. Thalhammer, O., Caradello, H., Pollak, A., Scheibenreiter, S., and Simbruner, G.: "Prospective and Retrospective Examination of an Easily Applicable Score to Predict the Probability of Premature Birth Defined by Weight", J. Perin. Med., 4:38, 1976.
26. Wilson, E.W. and Sill, H.K.: "Identification of the High Risk Pregnancy by a Scoring System", New Zealand Medical Journal, p. 437, November 28, 1973.
27. Haeri, A.D. and Naldrett, J.: "A Scoring System for Identifying Pregnant Patients with a High Risk of Perinatal Mortality", J. Obstet. Gynecol. Br. Common., 81:535, July 1974.
28. Fedrick, J.: "Antenatal Identification of Women at High Risk of Spontaneous Pre-Term Birth", Br. J. Obstet. Gynecol., 83:351, 1976.

29. Edwards, L.E., Barrada, M.E., Tatreau, R.W. and Hakanson, E.Y.: "A Simplified Antepartum Risk Scoring System", Obstet. Gynecol., 54(2):237, 1979.
30. Staisoh, K.J., Westlake, J.R. and Bashore, R.A.: "Blind Oxytooin Challenge Test and Perinatal Outcome", Am. J. Obstet. Gynecol., 138(4):399-403, October 15, 1980.
31. Pavelka, R., Riss, P., Parshalk, O. and Reinold, E.: "Practical Experiences in the Prevention of Prematurity Using Thalhammer's Score", J. Perinat. Med., 8(2):100-108, 1980.
32. Akhtar, J. and Sehgal, N.N.: "Prognostic Value of a Prepartum and Intrapartum Risk-Scoring Method", South Med. J., 73(4):411-414, April 1980.
33. Halliday, H.L., Jones, P.K. and Jones, S.L.: "Method of Screening Obstetric Patients to Prevent Reproductive Wastage", Obstet. Gynecol., 55(5):656-659, May 1980.
34. Cameron, R.: "Should Air Hostesses Continue Flight Duty During the First Trimester of Pregnancy?", Aerospace Medicine, 44(5): 552, May 1972.
35. Diddle, A.: "Gravid Women at Work: Fetal and Maternal Morbidity, Employment Policy and Medicolegal Aspects", Journal of Occupational Medicine, 12:10-15, 1970.
36. Guidelines on Pregnancy and Work (American College of Obstetricians and Gynecologists), U.S. Department of Health, Education and Welfare, Rockville, MD., DHE W(N10SH) Pub. No. 78-118, Sept. 1977.
37. Hendershot, G.E.: "Work During Pregnancy and Subsequent Hospitalization of Mothers and Infants", Public Health Reports, 94(5):425-31, Sept-Oct 1979.

38. Seitchik, J.: "Body Composition and Energy Expenditure During Rest and Work in Pregnancy", American Journal of Obstet. and Gynecol., 97(5):701-713, March 1967.
39. Tafari, N., et al.: "Effects of Maternal Undernutrition and Heavy Physical Work During Pregnancy on Birth Weight", British Journal of Obstet. and Gynecol., 87(3):222-27, March 1980.
40. Scholten, Paul: "Pregnant Stewardess - Should She Fly?" Aviation, Space and Environmental Medicine, 47(1):77-81, 1976.
41. Woodward, Susan L.: "How Does Strenuous Maternal Exercise Affect the Fetus? A Review", Birth and the Family Journal, 8:17, Spring, 1981.
42. Peckham, C.H.: "Uterine Bleeding During Pregnancy, Part 1: When Not Followed by Immediate Termination of Pregnancy", Obstet. Gynecol., 35(6):937, 1970.
43. Bjerre, B. and Bjerre, I.: "Significance of Obstetric Factors in Prognosis of Low Birthweight Children", Acta. Pediatr. Scand., 65:577, 1976.
44. McDonald, A.: "Retarded Foetal Growth", Clin. Develop. Med., 1965, p. 14.
45. Turnbull, E.P.N.: "The Outcome of Pregnancy Complicated by Threatened Abortion", J. Obstet. Gynecol., 1956:553.
46. Ornoy, A., Benady, S., Kohen-Raz, R., and Russell, A.: "Association Between Maternal Bleeding During Gestation and Congenital Anomalies in the Offspring", Am. J. Obstet. Gynecol., 124(5):474, 1976.
47. Asanti, R., and Tuulikki, V.: "Effect of Threatened Abortion on Foetal Prognosis", Acta. Obstet. et Gynecol. Scand., 42:107, 1963.

48. Funderburk, S.J., Guthrie, D., and Meldrum, D.: "Outcome of Pregnancies Complicated by Early Vaginal Bleeding", Br. J. Obstet. Gynecol., 87:100, 1980.
49. Butler, N., and Bonham, D.: "Perinatal Mortality: The First Report of the 1958 British Perinatal Mortality Survey", Edinburgh, 1963, p. 102, Edited by: E. and S. Livingstone.
50. Pasamanick, B., and Lilienfeld, A.M.: "Association of Maternal and Fetal Factors With Development of Mental Deficiency. Pt. 1: Abnormalities in the Prenatal and Paranatal Periods", J. Am. Med. Assoc., 159(3):155, 1955.
51. Lilienfeld, A.M., and Pasamanick, B.: "Association of Maternal and Fetal Factors With the Development of Epilepsy. Pt. 1: Abnormalities in the Prenatal and Paranatal Periods", J. Am. Med. Assoc., 155(8):719, 1954.
52. Fedrick, J., and Anderson, A.B.M.: "Factors Associated with Spontaneous Pre-Term Birth", Br. J. Obstet. Gynecol., 83:342, 1976.
53. Fedrick, J. and Adelstein, P.: "Factors Associated with Low Birth Weight of Infants Delivered at Term", Br. J. Obstet. Gynecol., 85(1):1, 1978.
54. Antonov, A.N.: "Children Born During the Siege of Leningrad in 1942", The Journal of Pediatrics, 30:250, 1947.
55. Bergner, L. and Susser, M.: "Low Birth Weight and Prenatal Nutrition: An Interpretative Review", Pediatrics, 46: 946, December 1980.
56. Butler, N.R. and Alberman, E.D. (eds.): "The Second Report of the 1958 British Perinatal Mortality Survey", Perinatal Problems, Edinburgh: E.S. Livingstone Limited, 1969.

57. Chalmers, Ian.: "The Search for Indices", Lancet, 2:1063, September 1979.
58. Edwards, L.E., et al: "Pregnancy in the Underweight Woman", Am. J. Obstet. Gynecol., 135:297, October 1, 1979.
59. Kessner, David M., et al: "Infant Death: An Analysis by Maternal Risk and Health Care", Washington: Institute of Medicine, 1973.
60. Leader, Arthur, et. al: "Maternal Nutrition in Pregnancy. Part 1: A Review", CMA Journal, 125:545, September 15, 1981.
61. Mora, J.L., et al: "Nutritional Supplementation and the Outcome of Pregnancy. 1. Birthweight", Am. J. Clin. Nutr., 32:455, 1979.
62. Mora, J.L., et al: "Nutritional Supplementation and the Outcome of Pregnancy. 3. Perinatal and Neonatal Mortality", Nutr. Rep. Int., 18:167, 1978.
63. Naismith, D.J.: "Maternal Nutrition and the Outcome of Pregnancy - A Critical Appraisal", Proc. Nutr. Soc., 39:1, 1980.
64. Ontario Department of Health. "Second Report of the Perinatal Mortality Study in Ten University Teaching Hospitals", Toronto, 1967.
65. Pencharz, Paul B.: "Maternal Nutrition and the Outcome of Pregnancy", CMA Journal, 125:529, September 15, 1981.
66. Rush, David, et al: "A Randomized Controlled Trial of Prenatal Nutritional Supplementation in New York City", Pediatrics, 65:683, April 1980.
67. Rush, David: "Nutritional Services During Pregnancy and Birthweight: A Retrospective Matched Pairs Analysis", CMA Journal, 125:567, September 15, 1981.

68. Singer, J., et al: "Relationship of Weight Gain During Pregnancy to Birth Weight and Infant Growth and Development in the First Year of Life", Obstet. Gynecol., 31:417, March 1968.
69. Smith, C.A.: "Effects of Maternal Undernutrition Upon the Newborn Infant in Holland", Journal of Pediatrics, 30:229, March 1947.
70. Susser, Mervin: "Prenatal Nutrition, Birthweight and Psychological Development: An Overview of Experiments, Quasiexperiments, and Natural Experiments in the Past Decade", Am. J. Clin. Nutr., 34: 784, April 1981.
71. U.S. Department of Health, Education and Welfare: "The Women and Their Pregnancies". The Collaborative Perinatal Study of the National Institute of Neurological Diseases and Stroke. Washington, D.C., 1972.
72. Warnyka, J., et al: "Healthiest Babies Possible. The Vancouver Project", The Canadian Nurse, 75:18, November 1979.
73. Davies, A. Michael: "Epidemiology of the hypertensive disorders of pregnancy", Bulletin of the World Health Organization, 57(3): 373-386, 1979.
74. Chesley, Leon C.: "The Remote Prognostic Significance of the Level of Blood Pressure in Pregnancy", Clinical and Experimental Hypertension, 777-801, 1980.
75. MacGillivray, I. and Campbell, Doris M.: "The Relevance of Hypertension and Oedema in Pregnancy", Clinical and Experimental Hypertension, 897-914, 1980.
76. Moodley, J.: "The Management of Hypertension in Pregnancy", S.A. Medical Journal, 58:103-109, 1980.
77. Fisher, K.A., Luger, A., Spargo, B.H. and Lindheimer, M.D.: "Hypertension in Pregnancy: Clinical-Pathological Correlations and Remote Prognosis", Medicine, 60(4):267-276, 1981.

78. Langford, H.G. and Watson, R.L.: "Prepregnant Blood Pressure, Hypertension During Pregnancy, and Later Blood Pressure of Mothers and Offspring", Hypertension, 2(4):130-133 (Suppl. 1), 1980.
79. Welt, S.I., Dorminy, J.H., Jelovsek, F.R., Crenshaw, M.C., and Gall, S.A.: "The Effect of Prophylactic Management and Therapeutics on Hypertensive Disease in Pregnancy: Preliminary Studies", Obstetrics and Gynecology, 57(5):557-565, 1981.
80. Curet, L.B., and Olson, R.W.: "Evaluation of a Program of Bed Rest in the Treatment of Chronic Hypertension in Pregnancy", Obstetrics and Gynecology, 53(3):336-340, 1979.
81. Gallery, E.D.M., Saunders, D.M., Hunyor, S.N. and Györy, A.Z.: "Randomised comparison of methyldopa and oxprenolol for treatment of hypertension in pregnancy", British Medical Journal, 1: 1591-1594, 1979.
82. Hauth, J.C., Cunningham, F.G. and Whalley, P.J.: "Management of Pregnancy-Induced Hypertension in the Nullipara", Obstetrics and Gynecology, 48(3):253-259, 1976.
83. Sawyer, M.M., Lipshitz, J., Anderson, G.D., Dilts, P.V. and Halperin, L.: "Diurnal and Short-Term Variation of Blood Pressure: Comparison of Preeclamptic, Chronic Hypertensive, and Normotensive Patients", Obstetrics and Gynecology, 58(3): 291-296, 1981.
84. Ounsted, M.K., Moar, V.A., Good, F.J. and Redman, C.W.G.: "Hypertension During Pregnancy With and Without Specific Treatment; the Development of the Children at the Age of Four Years", British Journal of Obstetrics and Gynecology, 87:19-24, 1980.
85. Dhall, K. and Dhall, G.I.: "Evaluation of the Supine Pressor Test", Gynecol. Obstet. Invest., 11:225-230, 1980.

86. Didolkar, S.M., Sampson, M.B., Johnson, W.L. and Petersen, L.P.: "Predictability of Gestational Hypertension", Obstetrics and Gynecology, 54(2):224-225, 1979.
87. Olmer, M., Gugliotta, J., Exbrayat, Ch., Gamberre, M., Grand-Vuillemin, M., et Ruf, H.: "Les accidents vasculo-renaux de la grossesse: étude de 75 cas évoluant chez des femmes indemnes d'antécédent rénal ou d'hypertension artérielle", Société de néphrologie, p. 824, Octobre 1979.
88. Beaufile, M., Donsimoni, R., Uzan, S., Colau, J.C., Salat-Baroux, J., et Richet, G.: "Données récentes sur la physiopathologie et la nosologie de l'hypertension artérielle gravidique", J. Urol. et de Nephrol., 12:805, 1979.
89. Equiedt, Ph., Tacquet, A., Puech, F., Cotteel, M., Potier, A., Leroy, J.L., et Delecour, M.: "Valeur pronostique de l'hyperuricémie dans les hypertensions artérielles de la grossesse", J. Gyn. Obst. Biol. Repr., 8:115, 1979.
90. Lambotte, R.: "Notions épidémiologiques récentes", J. Gyn. Obst. Biol. Repr., 9(1):33, 1980.
91. Vokaer, A.: "Facteurs participant éventuellement au déterminisme de l'hypertension gravidique. 1) La Régulation hormonale de bilan hydro-minéral. Perturbations de la balance hydro-minérale", J. Gyn. Obst. Biol. Repr., 9(1):39, 1980.
92. Fournier, A., et Pontonnier, G.: "Vascularisation placentaire et rénale en rapport avec des facteurs humoraux ou hormonaux locaux", J. Gyn. Obst. Biol. Repr., 9(1):45, 1980.
93. Lambotte, R., et Lecomte, J.: "Hypothèses interprétatives", J. Gyn. Obst. Biol. Repr., 9(1):53, 1980.
94. Ulstein, M., Sagen, N. and Eikhom, S.N.: "A Comparative Study of Labour Induced by Oral Prostaglandin E₂ and Buccal Tablets of Demoxytocin", Int. J. Gynaecol. Obstet., 17:243-245, 1979.

95. Piccolo, A.R.: "Clinical management of postdate pregnancy in the small community hospital", Journal of A.O.A. 80(4):271-275, 1980.
96. Freeman, R.K., et al: "Postdate pregnancy: Utilization of contraction stress testing for primary fetal surveillance", Am. J. Obstet. Gynecol., 140(2):128-135, 1981.
97. Gauthier, R.J., Griego, B.D. and Goebelsmann, U.: "Estriol in pregnancy", Am. J. Obstet. Gynecol., 139(4):382-389, 1981.
98. Homburg, R., Ludomirski, A. and Insler, V.: "Detection of Fetal Risk in Postmaturity", British J. of Obstet. Gynaecol., 86:759-764, 1979.
99. Callenbach, J.C. and Hall, R.T.: "Morbidity and Mortality of Advanced Gestational Age: Post-Term or Postmature", Obstetrics and Gynecology, 53(6):721-724, 1979.
100. Knox, G.E., et al: "Management of prolonged pregnancy: Results of a prospective randomized trial", Am. J. Obstet. Gynecol. 134(4):376-381, 1979.
101. Hauth, J.C., et al: "Post-Term Pregnancy. I.", Obstetrics and Gynecology, 56(4):467-470, 1980.
102. Green, J.N. and Paul, R.H.: "The Value of Amniocentesis in Prolonged Pregnancy", Obstetrics and Gynecology, 51(3):293-298, 1977.
103. Chambers, L.W. and West, A.E.: "The St. John's Randomised Trial of the Family Practice Nurse: Health Outcomes of Patients", Int. J. Epid., 7:153-160, 1979.
104. Eastman, N.J.: "Prematurity from the Point of View of the Obstetrician", Am. Prac., 1:343-352, 1947.
105. Gortmaker, S.L.: "The Effects of Prenatal Care on the Health of the Newborn", A.J.P.H., 69:653-660, 1979.

106. Institute of Medicine, National Academy of Sciences: "Infant Death, An Analysis by Maternal Risk and Health Care", In Contrasts in Health Status, Vol. 1, 1973.
107. Levy, B.S., Wilkinson, F.S. and Marine, W.M.: "Reducing Neonatal Mortality Rate with Nurse Midwives", Am. J. Obstet. Gynecol., 109:50-58, 1971.
108. Oppenheimer, E.: "Population Changes and Perinatal Mortality", A.J.P.H., 51:208-216, 1961.
109. Pakter, J., Rosner, H.J., Jacobziner, H. and Greenstein, F.: "Out of Wedlock Births in New York City. 2. Medical Aspects", A.J.P.H., 51:846-865, 1961.
110. Quick, J.D., Greenlick, M.R., and Roghmann, K.J.: "Prenatal Care and Pregnancy Outcome in an H.M.O. and General Population: A Multivariate Cohort Analysis", A.J.P.H., 71:381-390, 1979.
111. Reid, M.L. and Morris, J.B.: "Perinatal Care and Cost Effectiveness: Changes in Health Expenditures and Birth Outcome Following the Establishment of a Nurse Midwife Program", Med. Care, 17:491-500, 1979.
112. Rosenberg, S.N., Albertson, P.C., Jones, E.E. and Roberts, R.S.: "Complications of Labour and Delivery Following Uncomplicated Pregnancy", Med. Care, 19:68-79, 1981.
113. Spitzer, W.O., et al.: "The Burlington Randomised Trial of the Nurse Practitioner", N.E.J.M., 290:251-256, 1974.
114. Terris, M. and Glasser, M.: "A Life Table Analysis of the Relation of Prenatal Care to Prematurity", A.J.P.H., 64:869-875, 1974.
115. Wiener, G. and Milton, T.: "Demographic Correlates of Low Birth Weight", Am. J. Epid., 91:260-272, 1970.

116. Ashbury, J.: "The Crisis of Childbirth: Can Information and Childbirth Education Help?", Journal of Psychosomatic Research, 24:9, 1980.
117. Beck, N.C. et al.: "The Prediction of Pregnancy Outcome: Maternal Preparation, Anxiety and Attitudinal Sets", Journal of Psychosomatic Research, 24:343, 1980.
118. Charles, A.G., et al "Obstetric and Psychological Effects of Psychoprophylactic Preparation for Childbirth", AJOG, 131: 44-52, May 1, 1978.
119. Cogan, R.: "Effects of Childbirth Preparation", Clinical Obstetrics and Gynecology, 23(1):1, March 1980.
120. Dalzell, I.: "Evaluation of a Prenatal Teaching Program", Nursing Research, 14:160-3, 1965.
121. Davis, C.D. and Morrone, F.A.: "An Objective Evaluation of a Prepared Childbirth Program", AJOG, 84:1196-1206, 1962.
122. Doering, S.G. and Entwisle, D.R.: "Preparation During Pregnancy and Ability to Cope with Labour and Delivery", American Journal of Orthopsychiatry, 45:825-837, October 1975.
123. Enkin, M.W., et al.: "An Adequately Controlled Study of the Effectiveness of PPM Training", In Morris, N. (ed) Psychosomatic Medicine in Obstetrics and Gynecology Proceedings of Third International Congress, London 1971, Basel, S. Karger: 1972, p. 62.
124. Hott, J.R.: "Pre- and Post-partum Comparison of Self and Spouse in Primiparous Lamaze Couples Who Share Delivery and Those Who Do Not", Nursing Research, 29(1):2026, Jan-Feb 1980.
125. Hughey, M.J., et al.: "Maternal and Fetal Outcome of Lamaze Prepared Patients", Obstetrics and Gynecology, 51(6):643-647, 1978.

126. Humenick, Sharron: "Mastery: The Key to Childbirth Satisfaction - A Review", Birth and the Family Journal, 8:79, Summer 1981.
127. Humenick, Sharron: "Mastery: The Key to Childbirth Satisfaction - A Study", Birth and the Family Journal, 8:84, Summer 1981.
128. Huttel, F.A., et al.: "A Quantitative Evaluation of Psychoprophylaxis in Childbirth", Journal of Psychosomatic Research, 16:81-92, April 1972.
129. Laird, M.D. and Hogan, M.: "An Elective Program on Preparation for Childbirth at the Sloane Hospital for Women", AJOG, 72:641-7, 1956.
130. Melzack, Ronald et al.: "Labour is still Painful after Prepared Childbirth Training", CMA Journal, 125:357, August 15, 1981.
131. Scott, J.R. and Rose, N.B.: "Effect of Psychoprophylaxis (Lamaze Preparation) on Labour and Delivery in Primiparas", New England Journal of Medicine, 294:1205-1207, 1976.
132. Thomas, H. and Wyatt, R.H.: "One Thousand Consecutive Deliveries Under a Training for Childbirth Program", AJOG, 61:205-9, 1951.
133. Thordarson, L. and Constanzo, G.: "An Evaluation of an Educational Program for Expectant Parents", Canadian Journal of Public Health, 67:117-21, 1976.
134. Timm, Margaret M.: "Prenatal Education Evaluation", Nursing Research, 28(6):338-342, Nov-Dec 1979.
135. Van Auken, W.B.D. and Tomlinson, D.R.: "An Appraisal of Patient Training for Childbirth", AJOG, 66:100-5, 1953.
136. Willmuth, R., et al.: "Satisfaction with Prepared Childbirth and Locus of Control", JOGN Nursing, 7:33-37, May-June 1978.
137. Zax, M.: "Childbirth Education, Maternal Attitudes and Delivery", AJOG, 123:185, 1975.

138. Deter, R.L., Harrist, R.B., Hadlock, F.P.: "The use of Ultrasound in the assessment of normal fetal growth: A Review", J. Clin. Ultrasound, 9(9):481, 1981.
139. Nelson, L.H.: "Comparison of methods for determining crown-rump measurement by real-time ultrasound", J. Clin. Ultrasound, 9:67, 1981.
140. Parker, A.J., et al.: "Reproducibility of fetal crown-rump length measurements obtained with real-time ultrasound systems compared with those of a conventional B-scanner", Br. J. Obstet. Gynecol., 88(7):734, 1981.
141. Ott, W.J.: "Clinical application of Fetal Weight determination by real time ultrasound measurements", Obstet. Gynecol., 57(6): 758, 1981.
142. Deter, R.L., Hadlock, F.P., Harrist, R.B., et al.: "Evaluation of three methods for obtaining fetal weight estimates using dynamic image ultrasound", J. Clin. Ultrasound, 9(9):421, 1981.
143. Farrant, P.: "Ultrasound measurement of fetal limb length", Br. J. Radiology, 54(644):660, 1981.
144. Neilson, J.P., Hood, V.D.: "Ultrasound in obstetrics and gynecology - Recent Developments", Brit. Med. Bull., 36(3):249, 1980.
145. Queenan, J.F., O'Brien, G.D., Campbell, S.: "Ultrasound measurement of fetal limb bones", Am. J. Obstet. Gynecol., 138:297, 1980.
146. Martinez, D.A., Barton, J.I.: "Estimation of fetal body and fetal head volumes", Am. J. Obstet. Gynecol., 137:78, 1980.
147. Jordaan, H.V.T., Clark, W.B.: "Prenatal determination of fetal brain and somatic weight by ultrasound", Am. J. Obstet. Gynecol., 136:54, 1980.

148. Ott, W.J., Callaghan, G.M., Ritter, H.A.: "Routine ultrasound in the low risk obstetrical population", J. Reprod. Med., 24:203, 1980.
149. Poll, V., Kasby, C.B.: "An improved method of fetal weight estimation using ultrasound measurements of fetal abdominal circumference", Br. J. Obstet. Gynecol., 86:922, 1979.
150. Adam, A.H., Robinson, H.P., Dunlop, C.: "A comparison of crown-rump length measurements using a real-time scanner in an antenatal clinic and a conventional B-scanner", Br. J. Obstet. Gynecol., 86:521, 1979.
151. Fox, H.E., Hohler, C.W.: "Fetal evaluation by real-time imaging", Clin. Obstet. Gynecol., 20:339, 1979.
152. Sabbagha, R.E.: "Ultrasound in High Risk Obstetrics" in current concepts in Obstetrics and Gynecology (SUSPAN) 1979.
153. Morgan, C.L., Trought, W.S., et al.: "Applications of real-time ultrasound in Obstetrics", J. Clin. Ultrasound, 7:108, 1979.
154. Pedersen, J.F.: "Ultrasound evidence of sexual difference in fetal size in first trimester", Br. J. Radiology, 54(642):546.
155. Deter, R.L., Harrist, R.B., Hadlock, F.P., et al.: "The use of ultrasound in the detection of Intrauterine Growth Retardation: A Review", J. Clin. Ultrasound, 10:9, 1982.
156. Neilson, J.P.: "Detection of the small-for-dates twin fetus by Ultrasound", Br. J. Obstet. and Gynecol., 88:27, 1981.
157. Wladimiroff, J.W., Laar, J.: "Ultrasonic measurement of fetal body size: A RCT", ACTA Obstet. Gynecol. Scand., 59:177, 1980.
158. Kurjak, A., Kirkinen, P., Latin, V.: "Biometric and dynamic ultrasound assessment of small-dates infants", Obstet. Gynecol., 56:281, 1980.

159. Neilson, J.P., Whitfield, C.K., et al.: "Screening for the small-for-dates fetus: a two-stage ultrasonic examination schedule", Br. Med. J., 280:1203, 1980.
160. Arabin, B., Hoffbauer, H., Pachaly, J.: "Control of fetal size with multiple ultrasonic measurements", Journal of Perinatal Medicine, (suppl.), 1:135-137, 1981.
161. Varma, T.R., Taylor, H., Bridges, C.: "Ultrasound assessment of fetal growth", Br. J. Obstet. Gynecol., 86:623, 1979.
162. Crane, J.P., Kapta, M.M.: "Prediction of intrauterine growth retardation via ultrasonically measured head/abdominal circumference ratios", Obstet. Gynecol., 54:497, 1979.
163. Daikoku, N.H., Johnson, J.W.C., Graf, C.: "Patterns of intrauterine growth retardation", Obstet. Gynecol., 54:211, 1979.
164. Wittman, B.K., Robinson, H.P., Artchison, T.: "The value of diagnostic ultrasound as a screening test for IUGR", Am. J. Obstet. Gynecol., 134:30, 1979.
165. Weinstein, H., Spirt, B.A.: "Sonographic diagnosis of intrauterine growth retardation in a dichomonic diamniotic twin", J. Clin. Ultrasound, 7:219, 1979.
166. Sabbagha, R.E.: "Intrauterine growth retardation: Antenatal diagnosis by ultrasound", Obstet. Gynecol., 52:252, 1978.
167. Wladimiroff, J.W., Byoemsma, C.A., et al.: "Ultrasonic assessment of fetal head and body sizes in relation to normal and retarded fetal growth", Am. J. Obstet. Gynecol., 131:857, 1978.
168. Persson, P.H., Grennert, L., Gennser, G.: "Diagnosis of intrauterine growth retardation by serial ultrasonic cephalometry", ACTA Obstet. Gynecol. Scand. (Suppl), 78:40, 1978.
169. Persson, P.H., Marsal, K.: "Monitoring of fetuses with retarded BPD growth", ACTA Obstet. Gynecol. Scand. (Suppl), 78:49, 1978.

170. Campbell, S., Thoms, A.: "Ultrasound measurement of the fetal head to abdomen circumference ratio in assessment of growth retardation", Br. J. Obstet. Gynecol., 84:165, 1977.
171. Wladimiroff, J.W., Bloemsa, et al.: "Ultrasonic assessment of fetal growth", ACTA Obstet. Gynecol. Scand., 56:37, 1977.
172. Crane, J.P., Kopta, M.M., et al.: "Abnormal fetal growth patterns: Ultrasonic diagnosis and management", Obstet. Gynecol., 50:205, 1977.
173. Arias, R.: "The diagnosis and management of intrauterine growth retardation", Obstet. Gynecol., 49:293, 1977.
174. Whetham, J.C.G., Muggah, H., Davids, S.: "Assessment of intra-uterine growth retardation by diagnostic ultrasound", Am. J. Obstet. Gynecol., 125:577, 1976.
175. Queenan, J.F., Kibargh, S.T., et al.: "Diagnostic ultrasound for the detection of intrauterine growth retardation", Am. J. Obstet. Gynecol., 129:856, 1976.
176. Sabbagha, R.E.: "Ultrasonic Evaluation of Fetal Congenital Anomalies", Clinics in Obstetrics and Gynecology, Vol. 7:1, April 1980.
177. Hobbins, J.C., Grannum, Pat, Berkowitz, R.C.: "Ultrasound in the diagnosis of congenital anomalies", Am. J. Obstet. Gynecol., 134:331, 1979.
178. Farrant, P.: "Antenatal diagnosis of duodenal atresia", Br. J. Radiology, 54(643):633, 1981.
179. Crutard, E.: "Ultrasonography before abortion", JAMA, 246(10): 1077, 1981.
180. Morrison, J., Brunello, L.P.: "The antenatal diagnosis of hyperplacental, hydramnios and fetal ascites", J. Clin. Ultrasound, 5:338, 1977.

181. Hadlock, F.P.: "Sonography of fetal urinary tract anomalies",
Am. J. Radiology, 137(2):261, 1981.
182. Ruvinsky, E.D., et al.: "In utero diagnosis of umbilical cord
hematoma by ultrasonography", Am. J. Ob. Gyn., 140(7):833,
1981.
183. Yaghoobian, J., et al.: "Antenatal diagnosis of encephalocele by ultrasound",
J. Reprod. Med., 26(5):274, 1981.
184. Sabbagha, R.E.: "Ultrasonic evaluation of fetal congenital
anomalies", Clinics in Obstet. Gynecol., 7, 1980.
185. Dawes, G.S.: "Numerical analysis of human fetal H.R.: the quality
of ultrasound records", Am. J. Obstet. Gynecol., 141(1):43, 1981.
186. Bovicelli, L., Orseni, L.F., Rizzo, N., et al "Estimation of
gestational age during the first trimester using real-time
measurements of fetal crown-rump length and biparietal diameter",
J. Clin. Ultrasound, 9:71, 1981.
187. Kurtz, A.B., Wapner, R.J., Kunz, R.J., et al.: "Analysis of
biparietal diameter as an accurate indication of gestation age",
J. Clin. Ultrasound, 8:319, 1980.
188. Sabbacha, R.E., Highey, M.: "Standardisation of sonar cephalometry
and gestational age", Obstet. Gynecol., 52:402, 1978.
189. Gonzalez, A.C., Dale, E., Byers, R.H.: "Limitations in predictors
of gestational age and birth weight by ultrasonic methods",
J. Clin. Ultrasound, 6:233, 1978.
190. Wiener, S.N., Flynn, M.J., et al.: "A composite curve of ultrasonic
biparietal diameters for estimating gestational age", Radiology,
122:781, 1977.
191. Warsof, S.T., Gohari, P., Berkowitz, D.L.: "The estimation of
fetal weight by computer assisted analysis", Am. J. Obstet.
Gynecol., 128:881, 1977.

192. Drumm, J.E., Clinich, J., Mackenzie, G.: "The ultrasound measurement of fetal crown-rump length as a method of assessing gestational age", Br. J. Obstet. Gynecol., 83:417, 1976.
193. Kurjack, A., Cecuk, S., Breyer, B.: "Predication of maternity in first trimester of pregnancy by ultrasonic measurement of fetal crown-rump length", J. Clin. Ultrasound, 4:83, 1976.
194. Morrison, J., McLennan, M.J.: "The theory, feasibility, and accuracy of an ultrasonic method of estimating fetal weight", Br. J. Obstet. Gynecol., 83:833, 1976.
195. Dwers, W.A., Hemsell, D.L.: "The use of ultrasound in multiple gestation", Obstet. Gynecol., 53:500, 1979.
196. Duncan, S.L.B., Ginz, B., et al.: "Use of ultrasound and hormone assays in the diagnosis management of twin pregnancy", Obstet. Gynecol., 53:367: 1979.
197. Haney, A.F., et al.: "Significance of biparietal diameter differences between twins", Obstet. Gynecol., 51:609, 1978.
198. Houlton, M.C.C.: "Divergent biparietal diameter growth rates in twin pregnancies", Obstet. Gynecol., 49:542, 1977.
199. Fulton, I.C.: "Accuracy of measurements of crown-rump length and BPD made by inexperienced operators using a real-time scanner", Br. Med. J., 282(6281):2011, 1981.
200. Duncan, A.W.: "Fetal measurements made by inexperienced operators using real-time scanner", Br. Med. J., 283(6289):503, 1981.
201. Van Dongen, L.: "Training the obstetrician in ultrasonography", S.A. Med. J., 60(9):355, 1981.
202. Langendoerfer, S., Haverkamp, A.D., Murphy: "Pediatrics follow-up of a randomized controlled trial of intrapartum fetal monitoring techniques", Journal of Pediatrics, 97(1):103-107, 1980.

203. Haverkamp, A.D., Orleans, M., et al.: "A Controlled Trial of the Differential Effects of Intrapartum Fetal Monitoring", American Journal of Obstetrics and Gynecology, 134:399, 1979.
204. Kelso, I., Parsons, J.: "An assessment of continuous Fetal Heart Rate monitoring in labour: A randomized clinical trial", American Journal of Obstetrics and Gynecology, 131(5):526-532, 1976.
205. Trimbos, J.B.; Keirse, M.J.: "Assessment of Ante-partum Cardio-Tocograms in High Risk Pregnancy", British Journal of Obstetrics and Gynecology, 87(4):261-269, April 1980.
206. Renou, P., Chang, J., Anderson, I., Wood, C.: "Controlled trial of fetal intensive care", American Journal of Obstetrics and Gynecology, 126(4):470, 1976.
207. Haverkamp, A.D., Thompson, H.E., McFee, J.G., Cetrulo, C.: "The evaluation of continuous fetal heart rate monitoring in high risk pregnancy", American Journal of Obstetrics and Gynecology, 125:310, 1976.
208. Keane, M.W.D., Horger, E.O., Vice, L.: "Comparative study of stressed and non-stressed antepartum fetal heart rate testing", Obstetrics and Gynecology, 57(3):320-4, March 1981.
209. Krebs, H.B., Petres, R.E., Dunn, L.J., Segretti, A.: "Intrapartum fetal heart monitoring", American Journal of Obstetrics and Gynecology, 138(2):213-9, September 1980.
210. Paul, R.H., Hon, E.H.: "Clinical fetal monitoring. V. Effect on perinatal outcome", American Journal of Obstetrics and Gynecology, 118(4):529-33, February 1974.
211. Westgren, M., Ingemarsson, E., Ingemarsson, I., Solum, T.: "Intrapartum Electronic Fetal Monitoring in Low Risk Patients", Obstetrics and Gynecology, 56(3):301-304, September 1980.

212. Phillips, W.D., Towell, M.E.: "Abnormal Fetal Heart Rate Associated with Congenital Abnormalities", British Journal of Obstetrics and Gynecology, 87(4):270-274, April 1980.
213. Liu, D.T.Y., Blackwell, R.J., Tukel, S.: "The Relevance of (antenatal) and Intrapartum Fetal Heart Rate Patterns to Fetal Outcomes", British Journal of Obstetrics and Gynecology, 85:270-277, 1978.
214. Goodlin, R.C., Haesslin, H.C.: "When is it Fetal Distress?", American Journal of Obstetrics and Gynecology, 128(4):440-447, 1977.
215. Nochimson, D.J., Cetrulo, C.L.: "Intrapartum Fetal Monitoring", Journal of Family Practice, 4(5):845, 1977.
216. Amato, J.C.: "Fetal Monitoring in a Community Hospital", Obstetrics and Gynecology, 50:269-274, 1977.
217. Lee, W.K., Baggish, M.S.: "The Effect of Unselected Intrapartum Fetal Monitoring", Obstetrics and Gynecology, 37(5):516-520, 1976.
218. Chik, L., Sokol, R.J., Rosen, M.G.: "Prediction of the One Minute APGAR Score from Fetal Heart Rate Data", Obstetrics and Gynecology, 48(4):452-455, 1976.
219. Tejani, N., Mann, L.I.: "Correlation of Fetal Heart Rate Patterns and Fetal pH with Neonatal Outcome", Obstetrics and Gynecology, 48(4):460-63, 1976.
220. Tintera, G., Newman, R.L.: "Fetal Monitoring: Its Effect on Perinatal Mortality and Cesarean Section and its Complications", American Journal of Obstetrics and Gynecology, 122:750-54, 1975.
221. Edington, P.T., Sibanda, J., Beard, R.W.: "Influence on Clinical Practice of Routine Intrapartum Fetal Monitoring", British Medical Journal, 3:341-3, 1975.

222. Shenker, L., Post, R.C., Seiler, J.S.: "Routine Electronic Monitoring of Fetal Heart Rate and Uterine Activity During Labour", Obstetrics and Gynecology, 46(2):185-89, 1975.
223. Thomas, G.: "The Aetiology, Characteristics, and Diagnostic Relevance of Late Deceleration Patterns in Routine Obstetric Practice", British Journal of Obstetrics and Gynecology, 82:121, 1975.
224. Sturbois, G., Tournaire, M., Breart, G.: "Prediction of one minute APGAR Score by Fetal Heart Rate Analysis: A Comparison Between Three Methods", International Journal of Gynecology and Obstetrics, 13:118-122, 1975.
225. Gabert, H.A., Stenchever, M.A.: "Electronic Fetal Monitoring as a Routine Practice in an Obstetrical Service: A Progress Report", American Journal of Obstetrics and Gynecology, 118(4):534-7, 1974.
226. Kelly, V.C., Kulkarni, D.: "Experiences with Fetal Monitoring in a Community Hospital", Obstetrics and Gynecology, 41(6):818-824, 1973.
227. Gabert, H.A., Stenchever, M.A.: "Continuous Electronic Monitoring of Fetal Heart Rate during Labour", American Journal of Obstetrics and Gynecology, 115(7):919-923, 1973.
228. Shenker, L.: "Clinical Experiences with Fetal Heart Rate Monitoring of 1000 Patients in Labour", American Journal of Obstetrics and Gynecology, 115(8):1111-6, 1973.
229. Schiffrin, B.S., Dame, L.: "Fetal Heart Rate Pattern", Journal of American Medical Association, 219(10):1322-5, 1972.
230. Paul, R.H.: "Clinical Fetal Monitoring", American Journal of Obstetrics and Gynecology, 113(5):573-77, 1972.

231. Paul, R.H., Hon, E.H.: "A Clinical Fetal Monitor", Obstetrics and Gynecology, 35(2):161-9, 1970.
232. Dixon, A.S.: "The Electronic Fetal Monitor: Should Every Mother Have One?", Canadian Family Physician, 27:1023-28, June 1981.
233. Hill, L.M.: "Diagnosis and Management of Fetal Distress", Mayo Clinic Proceedings, 54(12):784-93, December 1979.
234. Hobbins, J.C., Freeman, R., Queenan, J.T.: "The Fetal Monitoring Debate", Obstetrics and Gynecology, 54(1):103-109, July 1979.
235. Goodlin, R.C.: "History of Fetal Monitoring", American Journal of Obstetrics and Gynecology, 133(3):323-52, February 1979.
236. Banta, H.D., Thacker, S.B.: "Assessing the Costs and Benefits of Electronic Fetal Monitoring", Obstetrics and Gynecology Survey, 34(8):627-42, 1979.
237. Editorial: "Intrapartum Fetal Monitoring for All?", British Medical Journal, II:1466, 1976.
238. Editorial: Obstetrical Gynecological Survey, 30:503-504, 1975.
239. Zuspan, F.P., Quilligan, E.J., Iams, J.D., Van Geijn, H.P.: "NICHD Consensus Development Task Force Report: Predictors of Intrapartum Fetal Distress - The Role of Electronic Fetal Monitoring", Editorial, American Journal of Obstetrics and Gynecology, 135(3):287-91, October 1979.
240. Banta, H.D., Thacker, S.B.: "Assessing the Costs and Benefits of Electronic Fetal Monitoring", Obstetrical and Gynecological Survey, 34(8):627-42, 1979.
241. Neutra, R.R., Fienberg, S.E., et al.: "Effect of Fetal Monitoring on Neonatal Death Rates", New England Journal of Medicine, 299:324-26, 1978.

242. Quilligan, E.J., Paul, R.H.: "Fetal Monitoring - Is it Worth it?", Obstetrics and Gynecology, 45(1):96-100, January 1975.
243. Starkman, M.N.: "Fetal Monitoring: Psychological Consequences and Management Recommendations", Obstetrics and Gynecology, 50:500-504, 1977.
244. Youngs, D.D., Starkman, M.N.: "Psychological and Physiological Aspects of Electronic Fetal Monitoring", Primary Care, 3:691-700, 1976.
245. Lewis, P.J., de Swiet, M., Baylan, P. and Bulpitt, C.J.: "How Obstetricians in the United Kingdom Manage Preterm Labour", B.J. Ob. Gyn., 87:574-77, July 1980.
246. Fuchs, F.: "Prevention of Prematurity", Am. J. Ob. Gyn., 126(7): 809-20, December 1, 1976.
247. Vital Statistics for Ontario: 1977.
248. Hull, D.: "Contribution of preterm birth to perinatal morbidity", In Preterm Labour. Edited by A. Anderson, R. Beard, M. Brudenell and P.M. Dunn, R.C.O.G., London, 1977.
249. Castren, O., Gummerus, M. and Sasrikoski, S.: "Treatment of Imminent Premature Labour", ACTA Obstet. Gyn. Scand., 54:95, 1975.
250. Csapo, A.I. and Herczeg, J.: "Arrest of Premature Labour by Isoxsuprine", Am. J. Ob. Gyn., 129(5):482-491, Nov. 1, 1977.
251. Henderson, P.A., Curet, L.B., Olson, R.W. and Schwartz, D.B.: "The Treatment of Premature Labor with Isoxsuprine", Wisc. Med. J., 79:29-33, Oct. 1980.
252. Wesselius-de Casparis, A., Thiery, M., Yolesian, A., Baumgarten, K., Brosens, I., Gamisans, O., Stolk, J.G. and Vivier, W.: "Results of Double-Blind, Multicentre Study with Ritodrine in Premature Labour", British Medical Journal, 17:144-147, July 1971.

253. Zlatnik, F.J. and Fuchs, F.: "A Controlled Study of Ethanol in Threatened Premature Labor", American Journal of Obstetrics and Gynecology, 112:610, 1972.
254. Ingemarsson, I.: "Effect of Terbutaline on Premature Labour: A Double-Blind Placebo-Controlled Study", American Journal of Obstetrics and Gynecology 125(4):520-24, June 15, 1976.
255. Laursen, N.H., Merkatz, I.R., Tejani, N., Roberson, A., Mann, L.I., Fuchs, F.: "Inhibition of Premature Labor: A Multi-center Comparison of Ritodrine and Ethanol", American Journal of Obstetrics and Gynecology, 127(8):837-845, April 15, 1977.
256. Steer, C.M., Petrie, R.H.: "A Comparison of Magnesium Sulfate and Alcohol for the Prevention of Premature Labor", American Journal of Obstetrics and Gynecology, 129(1):1-4, Sept. 1, 1977.
257. Richter, R.: "Evaluation of Success in Treatment of Threatening Premature Labor by Betamimetic Drugs", Am. J. Obstet. Gynecol., 127(5):482-486, March 1, 1977.
258. Richter, R., Hinselmann, M.J.: "The Treatment of Threatened Premature Labor by Betamimetic Drugs: A Comparison of Fenoterol and Ritodrine", Obstetrics and Gynecology, 53(1):81-87, January 1979.
259. Spellacy, W.N., Cruz, A.C., Birk, S.A. and Buti, W.C.: "Treatment of Premature Labor with Ritodrine: A Randomized Controlled Study", Obstetrics and Gynecology, 54(2):220-223, August 1979.
260. Creasy, R.K., Golbus, M.S., Laros, R.K., Parer, J.T., Roberts, J.M.: "Oral Ritodrine Maintenance in the Treatment of Premature Labor", American Journal of Obstetrics and Gynecology, 137(2):212-219, May 15, 1980.

261. Larsen, J.F., Hansen, M.K., Hesselake, H., Kristoffersen, K.,
Larsen, P.K., Osler, M., Weber, J., Eldon, K. and Lange, A.:
"Ritodrine in the Treatment of Preterm Labour: A Clinical
Trial to Compare a Standard Treatment With Three Regimens
Involving the Use of Ritodrine", British Journal of Obstetrics
and Gynecology, 87:949-957, November 1980.
262. Merkatz, I.R., Peter, J.B. and Barden, T.P.: "Ritodrine
Hydrochloride: A Betamimetic Agent for Use in Preterm Labor:
II Evidence of Efficacy", Obstetrics and Gynecology, 56(1):7-12,
July 1980.
263. Johnson, J.W.C., Austin, K.L., Jones, G.S., Davis, G.H. and
King, T.M.: New England Journal of Medicine, 293:675, 1975.
264. Watring, W.G., Benson, W.L., Wiebe, R.A. and Vaughn, D.L.:
"Intravenous Alcohol - A Single Blind Study in the Prevention
of Premature Delivery: A Preliminary Report", Journal of
Reproductive Medicine, 16:35, 1976.
265. Fuchs, F.: "Prevention of Prematurity", American Journal of
Obstetrics and Gynecology, 126:809, 1976.
266. Walters, W.A.W. and Wood, C.: "A Trial of Oral Ritodrine for
the Prevention of Premature Labour", British Journal of
Obstetrics and Gynecology, 84:26-30, January 1977.
267. Tambyraja, R.L., Atputharajah, V., Salmon, Y.: "Prevention of
Prematurity in Twins", Australia, New Zealand, Journal of
Obstetrics and Gynecology, 18:179-181, 1978.
268. O'Connor, M.C., Murphy, H. and Dalrymple, I.J.: "Double Blind
Trial of Ritodrine and Placebo in Twin Pregnancy", British
Journal of Obstetrics and Gynecology, 86:706-709, 1979.
269. Mehra, P., Raghvan, K.S., Devi, P.K. and Chaudhury, R.: "Effect
of Intravenous Alcohol on Premature Labour", International
Journal of Gynecology and Obstetrics, 8:160, 1970.

270. Graff, G.: "Failure to Prevent Premature Labor With Ethanol", American Journal of Obstetrics and Gynecology, 110:878, 1971.
271. Lewis, P.J. and Baylan, P.: "Alcohol and Fetal Breathing", Lancet, 1:388, 1979.
272. Guilliams, S. and Held, B.: "Contemporary Management and Conduct of Preterm Labor and Delivery: A Review", Ob. Gyn. Surv., 34(3): 248-255, 1979.
273. Sharpe, G.L., Thalme, B. and Larsson, K.S.: "Studies on the closure of ductus arteriosus, XI. Ductal closure in utero by prostaglandin synthetase inhibitors", Prostaglandins, 8:863, 1974.
274. Steer, C.M. and Petrie, R.H.: "A Comparison of Magnesium Sulfate and Alcohol for the Prevention of Premature Labor", Am. J. Ob. Gyn., 129(1):1-4, Sept. 1, 1977.
275. Barden, T.P., Peter, J.B. and Merkatz, I.R.: "Ritrodrene Hydrochloride: A Betamimetic Agent for Use in Preterm Labor", Obs. Gyn., 56(1):1-6, July 1980.
276. Zuckerman, H., Reiss, U. and Rubinstein, I.: "Inhibition of Human Premature Labor by Indomethacin", Obstetrics and Gynecology, 44: 787, 1974.
277. Schenken, R.S., Hayashi, R.H., Valenzuela, G.V., Castillo, M.S.: "Treatment of Premature Labor with beta sympathomimetics: Results with Isoxsuprine", American Journal of Obstetrics and Gynecology, pp. 773-780, August 1, 1980.
278. Wallace, R.L., Ansbacher, R. and Otterson, W.N.: "Inhibition of Premature Labor by Terbutaline", Obstetrics and Gynecology, 51(4):387-392, April 1978.

279. Stubblefield, P.G.: "Pulmonary Edema Occurring after Therapy with Dexamethasone and Terbutaline for Premature Labor: A Case Report", American Journal of Obstetrics and Gynecology, 132:341, 1978.
280. Elliott, H.R. and Abdulla, U.: "Pulmonary Oedema Associated With Ritodrine Infusion and Betamethasone Administration in Premature Labour", British Medical Journal, 16:799-800, September 1978.
281. Katz, M., Robertson, P.A. and Creasy, R.K.: "Cardiovascular Complications Associated with Terbutaline Treatment for Preterm Labor", American Journal of Obstetrics and Gynecology, pp. 605-608, March 1, 1981.
282. Lind, T., Godfrey, K.A., Gerrard, J. and Bryson, M.R.: "Continuous Salbutamol Infusion Over 17 Weeks to Pre-Empt Premature Labour", The Lancet, pp. 1165-1166, November 29, 1980.
283. Tamby Raja, R.L., Ng, S.C. and Ratnam, S.S.: "Fenoterol: An Effective Agent for the Postponement of Preterm Birth", New Zealand Medical Journal, 93(681):218-20, April 8, 1981.
284. Chimura, T.: "The Treatment of Premature Labor by Drugs", Nippon Sanka Fujinka Gakkai Zasshi, 32(10):1620-4, October 1980.
285. Chance, G.W., Matthew, J.D., Williams, G. and Cunningham, K.: "Neonatal Transport: A Controlled Study of Skilled Assistance", Pediatrics, Vol. 93(4):662-666, 1978.
286. Sumners, J., Harris, H.B., Jones, B., Cassidy, G. and Wirtschaffer, D.D.: "Regional Neonatal Transport: Impact of an Integrated Community Center System", Pediatrics, Vol. 65(5):910-916, 1980.
287. Cats, B.P. and Beganovic, N.: "Intensive Care for Sick Neonates, the Earlier the Better: Improved Survival Rate of IRDS Children, Referred by Means of a Mobile Neonatal Intensive Care Unit", J. Perinat. Med., 7:19-22, 1979.

288. Skelton, M.A., Perkett, E.A., Major, C.W., Vaughan, R.L. and Stahlman, M.T.: "Transport of the Neonate", South. Med. J., Vol. 72(2):144-148, Feb. 1979.
289. Blake, A.M., McIntosh, N., Reynolds, E.O.R. and Andrew, D. St.: "Transport of Newborn Infants for Intensive Care", Br. Med. J., pp. 13-17, October 4, 1975.
290. Cunningham, M.D. and Smith, F.R.: "Stabilization and Transport of Severely Ill Infants", Ped. Clin. N. Am., Vol. 20(2):359-367, May 1973.
291. Chance, G.W.: "Transportation of High Risk Newborn Infants: The Role of the Trained Nurse; Report and Recommendations", Ontario Ministry of Health Demonstration Model Grant #DM174.
292. Storrs, C.N. and Taylor, M.R.H.: "Transport of Sick Newborn Babies", Br. Med. J., pp. 328-332, August 8, 1970.
293. Ryan, M.E.: "Transport of Neonate for Intensive Care", J. Am. Ob. Assoc., Vol. 78:pp. 103/75-109/81.
294. Souma, M.L.: "Maternal Transport: Behind the Drama", Am. J. Obstet. Gynecol., 134(8):904-910, August 15, 1979.
295. Boehm, F.H. and Haire, M.F.: "One-way Maternal Transport: An Evolving Concept: Inpatient Services", Am. J. Ob. Gyn., 134(4): 484-492, June 15, 1979.
296. Knuppel, R.A., Cetrulo, C.L., Ingardia, G.J. et al.: "Experience of a Massachusetts Perinatal Centre", N. Eng. J. Med., 30(10):560-562, March 8, 1979.
297. Giles, H.R., Isaman, J., Moore, W.J., Christian, C.D.: "The Arizona High-risk Maternal Transport System: An Initial View", Am. J. Obstet. Gynecol., 128(4):400-407, June 15, 1977.
298. Harris, B.A., Wirtschaffer, D.D., Huddleston, J.F. and Perlis, W.: "In Utero Versus Neonatal Transportation of High-risk Perinates: A Comparison", Ob. Gyn., 57(4):496-499, April 1981.

299. Levy, D.L., Noelke, K., Goldsmith, J.P.: "Maternal and Infant Transport Program in Louisiana", Ob. Gyn., 57(4):500-504, April 1981.
300. Modanlou, H.D., Dorchester, W., Freeman, R.K. and Rommal, C.: "Perinatal Transport to a Regional Perinatal Center in a Metropolitan Area: Maternal versus Neonatal Transport", Am. J. Ob. Gyn., 138(8):1157-1164, December 15, 1980.
301. Modanlou, H.D., Dorchester, W.L., Thorosian, A. and Freeman, R.K.: "Antenatal Versus Neonatal Transport at a Regional Perinatal Center: A Comparison Between Matched Pairs", Ob. Gyn., 53(6): 725-729, June 1979.
302. Merenstein, G.B., Pettett, G., Woodall, J. and Hill, J.M.: "An Analysis of Air Transport Results in the Sick Newborn: Antenatal and Neonatal Referrals", Am. J. Obstet. Gynec., 128(5):520-524, July 1, 1977.
303. Harris, T.R., Isaman, J. and Giles, H.R.: "Improved Neonatal Survival Through Maternal Transport", Ob. Gyn., 52(3):294-300, Sept. 1978.
304. Bhagwanani, S.G., Fahmy, D. and Turnbull, A.C.: "Prediction of neonatal respiratory distress by estimation of amniotic fluid lecithin", Lancet, 1:159-162, 1972.
305. Bichler, A., Daxenbichler, G. and Geir, W.: "Determination of amniotic fluid palmitic acid concentration for the estimation of fetal lung maturity", Clin. Chim. Acta, 74:133-138, 1977.
306. Jackson, B.A., Anderson, G.D. and Meld, B.: "Amniotic fluid phospholipids and fetal lung maturity", Am. J. Obstet. Gynecol., 121:1095-1099, 1975.
307. Olson, E.B., Hartline, J.V., Schneider, J.M. and Groven, S.N.: "The use of amniotic fluid bubble stability, L/S ratio and creatinine concentration in the assessment of fetal maturity", Am. J. Obstet. Gynecol., 122:755-760, 1975.

308. Russell, P.T., Miller, W.J. and McLain, C.R.: "Palmitic acid content of AF lecithin as an index to fetal lung maturity", Clin. Chem., 20:1431-1434, 1974.
309. Stark, R.I., Blumenheld, F.A., Cheskin, W.S., Dynenfurth, I. and Jones, L.S.: "Amniotic fluid fluorescence polarisation value as a predictor of R.D.S.", J. Pediatrics, 96:301-304, 1980.
310. Warren, C., Holton, J.B. and Allen, J.T.: "Assessment of fetal lung maturity by estimation of AF palmitic acid", Brit. Med. J., 11:94-96, 1974.
311. Werner, M., Genta, V.M. and Williams, D.: "Optimization of diagnostic discrimination applied to the amniotic fluid lecithin/sphingomyelin ratio", Clin. Chim. Acta., 101:155-162, 1980.
312. Wilde, C.E. and Oakey, R.E.: "Biochemical tests for the assessment of feto-placental function", Am. Clin. Biochem., 12:83-118, 1975.
313. Browne, Helen E. and Isaacs, Gertrude: "The Frontier Nursing Service: The Primary Care Nurse in the Community Hospital", Am. J. Obstet. Gynecol., 124:14, January 1, 1976.
314. Burnett, John E.: "A Physician-Sponsored Community Nurse-Midwife Program", Obstetrics and Gynecology, 40:719, November 1972.
315. Cherry, Janet and Foster, Joyce Cameron: "Comparison of Hospital Charges Generated by Certified Nurse-Midwives' and Physicians' Clients", Journal of Nurse-Midwifery, 27:7, Jan-Feb 1982.
316. Dillon, Thomas F., et al.: "Midwifery, 1977", Am. J. Obstet. Gynec., 130:917, April 15, 1978.
317. Haire, Doris: "Improving the Outcome of Pregnancy Through Increased Utilization of Midwives", Journal of Nurse-Midwifery, 26:5, Jan-Feb 1981.

318. Hewitt, Margaret A. and Hangleben, Karin Larson: "Nurse-Midwives in a Hospital Birth Centre", Journal of Nurse-Midwifery, 26:21, Sept-Oct 1981.
319. Levy, Barry S., et al.: "Reducing Neonatal Mortality Rate with Nurse-Midwives", Am. J. Obstet. Gynec., 109:50, January 1, 1971.
320. Mann, Rosemary J.: "San Francisco General Hospital Nurse-Midwifery Practice: The First Thousand Births", Am. J. Obstet. Gynec., 140: 676, July 15, 1981.
321. Record, Jane Cassels and Cohen, Harold R.: "The Introduction of Midwifery in a Prepaid Group Practice", American Journal of Public Health, 354, March 1972.
322. Reid, Michael L. and Morris, Jeffrey B.: "Perinatal Care and Cost-Effectiveness", Medical Care, 17:491, May 1979.
323. Slome, C., et al.: "Effectiveness of Certified Nurse-Midwives", Am. J. Obstet. Gynecol., 124:177, January 15, 1976.

APPENDIX 2

TASK FORCE MEMBERS AND CONSULTANTS

APPENDIX 2

TASK FORCE MEMBERS AND CONSULTANTS

MEMBERS

Chance, Graham W: Birmingham (Eng)'56; MB, ChB; FRCP (Lond); DCH; Paed; FRCP(C); Neonatologist, Director of Nurseries, St. Joseph's Hospital, London, Ontario; Consulting Staff, War Memorial Children's Hospital; Professor, Department of Pediatrics and Department of Obstetrics and Gynaecology, University of Western Ontario; Co-ordinator, Regional Perinatal Program; Chairman, Canadian Coalition for Prevention of Handicap of Perinatal Origin.

Dunne, James T: University of Dalhousie 1964; FRCS(C), ObstGyn; Staff, St. Clare's Mercy and Grace General Hospitals and Health Sciences Centre; Chairman, Department of Obstetrics and Gynecology, Memorial University, St. John's, Newfoundland.

Gagnon, Martine: University of Montreal 1973; FRCS(C), ObstGyn; Department of Obstetrics and Gynecology, Maisonneuve-Rosemont Hospital, Montreal, P.Q.

Hamilton, Emily: University of McGill, MDCM 1975; LMCC 1975; FRCS(C), ObstGyn 1979; CSPQ 1979; Director of Obstetrical and Gynecology, Ultrasonography, Women's Pavilion, Royal Victoria Hospital; Assistant Obstetrician and Gynecologist, Royal Victoria Hospital; Clinical Assistant Obst. and Gyn., Jewish General Hospital; Assistant Professor, Department of Obstetrics and Gynecology, Department of Medicine, McGill University; Consultant Obstetrics and Gynecology, Herzl Family Practice Centre; Consultant in Obst. and Gyn., Baffin Zone, Northwest Territories.

Kaufman, Karen: R.N., Clinical Nurse Specialist, Obstetrics and Gynecology; Faculty of Health Sciences, McMaster University, Hamilton, Ontario.

Kaye, Joyce (MacDonald): R.N.; Coordinator, Nova Scotia Reproductive Care Program, Halifax, N.S.

Leduc, Bernard: University of Montreal, M.D. 1963 D. Phil; C.S.P.Q.; F.A.C.O.G.; Associate Professor, Obstetrics and Gynecology, University of Montreal; Chief, Perinatology, Hôpital Notre-Dame, Montreal.

Lennox, Robert H: McGill 1943; DTM; MPH; FRCP(C), paediatrician; Senior Medical Consultant, Health Services & Promotion Branch, Department of National Health and Welfare; Courtesy Staff, Children's Hospital of Eastern Ontario; *Secretary of Federal Task Force on High Risk Pregnancy.

MacLachlan, Thomas B, M.D.: University of Toronto 1952; FRCS(C), ObstGyn; Staff, University Hospital, Saskatoon, Saskatchewan; Professor and Head, Department of Obstetrics and Gynecology, University of Saskatchewan.

Martin, Thomas R: Durham (Eng)'59; MB, BS; FACOG; FRCS(C), ObstGyn; Chief, Department of Obstetrics and Gynecology, St. Paul's Hospital, Vancouver, British Columbia; Associate Professor, University of British Columbia.

Milligan, John E., M.D.: University of Toronto 1959; FACOG; FRCS(C), ObstGyn; Obstetric Director, Perinatal Unit, Women's College Hospital; Associate Professor, Departments of Obstetrics and Gynaecology and Pediatrics, University of Toronto; Chairman of Ontario Medical Association Committee on Perinatal Care; Research Perinatology and Fetal Physiology.

Milner, Ruth: B.Sc., M.I.S.; Assistant Professor, Department of Clinical Epidemiology and Biostatistics, McMaster University, Hamilton 1978; formerly attached to Medical Research Council Epidemiology and Tropical Metabolism Research Units, Jamaica, W.I.; Major Research in Perinatal Area of Medicine and Obstetric and Neonatal Studies including High Risk Pregnant Women and Low Birthweight Infants.

Morrison, Ian: Glasgow (Scotland)'59; MB, ChB; FRCS(C), ObstGyn; Staff, St. Boniface General Hospital, Winnipeg, Manitoba; Consultant Staff, Victoria Hospital; Assistant Professor, Obstetrics and Gynecology, University of Manitoba.

Peddle, Leo J., M.D.: University of Dalhousie 1962; FACOG; FRCS(C), ObstGyn; Professor, Department of Obstetrics and Gynecology and Chairman, Division of Perinatal Medicine, University of Dalhousie, Nova Scotia.

Post, Shirley: R.N., B.Sc. (Ed), University of Ottawa 1967, M.H.A., President, Association for Care of Children's Health; Executive Director, Canadian Institute of Child Health, Ottawa, Ontario.

Rae, Don W., M.D.: University of Manitoba 1957; CCFP; FCFP; R.C.G.P.; Staff, Portage District General Hospital, Portage LaPrairie; Assistant Professor, Family Medicine, University of Manitoba.

Smith, Douglas M.: Cambridge (Eng)'56; MB, BCh; MRCOG; FRCS(C), ObstGyn; Staff, University Hospital, University of Alberta; Courtesy Staff, Charles Cammell Hospital; Teaching Staff, University of Alberta.

Stewart, Hugh V., M.D.: University of Alberta 1970; MCFP; Association Staff, St. Michael's General and Lethbridge Municipal Hospitals.

Walters, Jack H., M.D.: University of Western Ontario 1951; MCFP(Hon); FACOG; FRCOG; DABOG; FRCS(C), ObstGyn; Professor, Department of Obstetrics and Gynecology, Faculty of Medicine, University of Ottawa; Obstetrics and Gynecology, Ottawa General Hospital. *Chairman of Federal Task Force on High Risk Pregnancy.

Members of Dr. Milner's group included the following:

John Lament, M.D., F.R.C.S.(C)
Neil Johnston, M.Sc.
Lynn Jones, R.N.
Carol Rand B.Sc.

CONSULTANTS

Banister, P.G.: Edinburgh (Scotland)'51; M.B., ChB; MPH (Calif); FRCP(C), Paed; Consultant Paediatrician, Health Services and Promotion Branch, Department of National Health and Welfare; Staff, Children's Hospital of Eastern Ontario.

Effer, S.B., M.D.: Toronto 1957; FACOG; FRCS(C), ObstGyn; Deputy Head, Department of Obstetrics, Grace Hospital, Vancouver, British Columbia 1982; Professor, Head, Division of Fetal Maternal Medicine, University of British Columbia 1981.

Fabia, J., M.D.: (Lille, France) 1944; Dip. Occupational Medicine (Paris, France) 1949; M.S. (Epid) Harvard, 1966; D.S. (Epid) Harvard (1970); Professor, Faculty of Medicine, Laval University, Quebec; National Health Research Scientist 1976-1982.

Murdoch, A.J.: M.D.C.M., McGill 1963; Ph.D. (London, England, 1974); FRCP(C) 1969; Associate Director General (West) Operations, Medical Services Branch, Department of National Health and Welfare, Ottawa.

Nelson, N.: B.A. (Manitoba) 1970; Ph.D. (Medical Sciences) 1979, McMaster University; Consultant (Epidemiology) Maternal and Child Health Coalition, Winnipeg, Manitoba; Co-ordinator (Vancouver) Health Promotion Services Program, Department of National Health and Welfare.

Swyer, Paul R.: Cambridge (Eng)'40; MA (Cantab); MRCP (Lond); DCH (Eng); FRCP(C), Paed.; Chief, Division of Perinatal Medicine and Senior Staff Physician, Hospital for Sick Children; Senior Scientist, Research Institute, Hospital for Sick Children; Professor, Department of Paediatrics, University of Toronto.

APPENDIX 3

REPORT OF SUBCOMMITTEE ON THE PRENATAL RECORD

APPENDIX 3

REPORT OF SUBCOMMITTEE ON THE PRENATAL RECORD

CONTENTS

1. Philosophy and Functions of Prenatal Record
2. Content of Prenatal Record
 - 2.1 Basic or Essential Information
 - 2.2 Optional Information
 - 2.3 Risk Factors
3. Design of Prenatal Form
4. Data Collection
5. Summary and Conclusions

1. Analysis of Provincial Prenatal Records and Their Functions

The recording and documentation of obstetric information has, over the years evolved into the collection of a relatively uniform data base contained in a prenatal record. This prenatal record is used in pregnancy as a single, complete and comprehensive summary of the several factors which might directly, or indirectly, influence the outcome of the pregnancy. At each visit the physician can quickly refresh his/her memory and evaluate any changes from previously recorded baselines. It also forms a record which can be transferred with a woman or used on the labour floor so that management at this time is appropriate to antecedent events. On reviewing the examples of the prenatal forms from eight provinces that were available to the Committee, these functions appear to be recognized in principle by all provincial records.

All provinces provide a mechanism for the physician to retain the original prenatal sheet while still allowing at least one copy of this information to be received by the hospital or the patient.

All provincial prenatal sheets recognize the need for certain basic background information and in addition provide line-by-line space for each prenatal visit in the current pregnancy. The items of information that are recorded at each prenatal visit are almost uniform throughout the country and the number of visits that can be recorded on one prenatal sheet varies from 14 (5 provinces) to 19 (Newfoundland). Each prenatal form lists a number of items as "reminders" (some check-off items) so that the record when completed will contain a consistent minimum data base for each pregnant patient. On the whole, there is surprisingly little variation in the basic content of the different prenatal sheets (see pages 94-95). Information related to history and ongoing prenatal visits is already relatively "standardized" in most instances.

Again, while the design format and number of pages contained within the record differs from province to province, it is quite

evident that each prenatal sheet from each province is an attempt to provide the physician with a practical document that is primarily used for patient management. The "reminder" or menu system, with or without a checklist, makes history taking and recording quicker, assists in a review of material at each subsequent woman's visit, and to some extent overcomes some of the problems of illegibility if more than one physician is involved in the woman's care. The predominant intent seems to be a functional record for use in patient management. None of the provincial forms appear to be specifically designed for the subsequent collection and storage of collective data or for computerization.

The principle of recognizing certain factors that may increase the risk of a poor maternal or fetal outcome for the current pregnancy, and of listing these as an aid to the attending physician, seems to have been established by at least seven of the eight provinces. Here, however, there is less uniformity. Some provinces use a broad grading or classification system in which there are differences in the choice of risk factors within each category. Others use a scoring system for risk identification and these numerical scoring systems differ between provinces.

In summary, it is apparent that all provincial prenatal forms have certain features in common. The standardization of the prenatal record as it has evolved in each province seems to be an attempt to assist the physician in the following:

1. Assure a complete basic record necessary for the proper management of the pregnant woman.
2. Reduce the amount of time taken to obtain this record.
3. To provide a means for the attending physician to review "at a glance" baseline data and information from previous prenatal visits which might influence subsequent management.

4. To provide some assistance to clinical judgment by indicating risk factors that allegedly influence outcome.
5. To provide a means of continuity for the recording of antecedent events that might influence management when the woman enters a labour unit or when the responsibility for patient care is transferred to another physician.

2. Content of Prenatal Forms

These functional recording systems are an attempt to meet the needs of both family physicians and specialists. It is important to recognize that compromises are necessary to achieve the needs of both groups. The proportion of deliveries carried out by family physicians varies from province to province but it is estimated that at least 50% of all deliveries in the country are still performed by this group. Therefore, in considering the function of a prenatal record, it is obvious that the needs of family physicians must be taken into consideration. Despite distinctive differences in referral patterns of obstetric practice between provinces and despite the need to serve both specialist and family physician there is in fact a fair degree of uniformity of content in all the prenatal forms. With minor adaptations each provincial prenatal form could contain the same basic information since differences that are currently existing are of a minor nature. An attempt at uniformity of content for specific high-risk factors may be a more contentious issue.

2.1 Factors considered basic or essential and common to most prenatal sheets

Method of patient identification
Identification of attending physician
Age, date of birth of patient
Parity
L.N.M.P.
E.D.C.
Menstrual cycle

Contraception/date discontinued

Genetic amniocentesis

Bleeding since L.N.M.P.

Past obstetric history

stillbirth, neonatal death, prematurity, abortion, number
of living children

Previous confinements

dates, birthweight, gender, gestation, type of delivery,
pregnancy/labour complications, neonatal complications

Past medical history

renal, cardiac, infertility, infections, diabetes, epilepsy,
urinary tract infections, allergies, previous blood trans-
fusions and operative procedures

Family history

diabetes, twins, congenital malformations, hereditary
deafness, T.B., hypertension

Present history

hyperemesis, febrile illnesses, current medications,
alcohol and drug use, smoking

Investigations

Hb, Rh, serology, Rubella, cytology, blood sugars, pregnancy
tests and mid-stream urine for culture

Initial examination

pre-pregnancy weight, height, current weight, complete
examination and pelvic examination, uterine size compared
to gestation

Repeat examinations or subsequent visits

dates, gestation, weight, urine analysis, blood pressure,
fundal height, presentation, fetal heart rate, remarks or
comments, time of next visit

Risk Identification

This process forms a component of most of the prenatal forms
that were reviewed. By common usage therefore it seems that

some form of risk identification process should be included as an essential component of the prenatal form. This is particularly the case if one of the purposes of the record is to assist family physicians; a list of risk factors may serve as a reminder for the particular patient under review.

2.2 Optional features for prenatal forms

These features may be of assistance to the attending physician but tend to be a matter of personal inclination. In addition, these items are not likely to significantly affect patient management. There is not a great deal of evidence that their documentation significantly affects the outcome of pregnancy. Some prenatal forms include space and reminders that summarize completed investigations. For example - Rh Gamma Globulin, ultrasound, G.T.T., medications, 28 weeks referral plan, consultation, etc.

Growth-curve charts are another feature as an aid to diagnosis of "large-for-dates" or "small-for-dates".

Documentation (optional)

Occupation of patient

Religion of patient

Ethnic origin of patient

Name of father

Occupation of father

Religion of father

Nutritional history

Hospital

Type of anesthesia requested

Labour requests

Father delivery

Rooming-in

Early discharge

Circumcision

Pediatrician

Discussion Topics (optional)

These appear to be a reminder of topics that could profitably be discussed with the patient; whether the majority of physicians have sufficient time to carry out an in-depth discussion of these topics may be another matter.

Weight gain

Smoking

Alcohol

Breast feeding

Prenatal classes

Work plans

Exercise

Coitus

2.3 Risk Factors

The validity of currently-used predictors of unfavourable outcome and the basis upon which these outcomes are identified is the topic of review by another sub-committee. The suggestion at this stage, in this document, is that some form of risk identification be retained either as an essential or optional feature of a prenatal record since most provinces have already taken this route. Provided they are used as an aid to clinical judgment, their inclusion probably serves a useful purpose particularly for practitioners who do not have a heavy obstetric caseload.

The Committee during a discussion on risk markers decided that a number were sufficiently obvious that further research into their validity was not necessary. Presumably the following list, agreed to by the major Committee, could form a minimum list:

Low birthweight

Previous stillbirth or neonatal death

Medical disorders
Rh Isoimmunization
Gestational diabetes
Multiple pregnancy
Premature spontaneous rupture of membranes
Prematurity
Malpresentation

Many other risk factors are included in the various prenatal forms that have been distributed to all Committee members. The inclusion or exclusion of these other risk factors will presumably depend on the outcome of the deliberations of the sub-committee reviewing this topic.

3. Design of Prenatal Form

In contrast to the content of the prenatal forms which contain many common basic medical/obstetric fields, the design of each provincial prenatal form varies considerably.

One point on design that could probably be agreed upon is that at least one copy of any prenatal form should be available so that a duplicate could be retained as part of the patient's hospital record. Improvements in design of a prenatal form should have input from those persons most closely involved in its use. This design should facilitate the functions of the prenatal record that have already been discussed.

4. Data Collection

Since each provincial prenatal form contains almost the same basic information with respect to each pregnancy it is tempting to consider the collection of data from this source. This is particularly the case with respect to information that might be obtained about the ratio of risk pregnancies handled by hospitals, the precise gestation in pregnancy that specific risks might be identified and the determination of any geographic differences in the proportion of risk

pregnancies. The prevalence of rubella susceptibility, nutritional status, or number of prenatal visits might be other examples of useful information on a provincial or national basis. However, there are certain difficulties connected with the use of prenatal forms for accurate data collection.

These are summarized below:

1. Compliance in the use of a specific form

Any attempt at data retrieval must capture nearly all of the obstetric population. This is impossible unless all physicians in a province use a standardized prenatal form.

2. Compliance for completion

A survey of over 2000 prenatal records by a Task Force of the Social Planning Council in Manitoba showed a wide range of variation on an item-per-item basis as far as completion rates were concerned. Some items had a completion rate of greater than 80%, others less than 50%. Obviously data collection with this type of variation will produce inaccuracies if there are a number of missing fields.

3. Accuracy of recorded data

In all prenatal sheets there are a number of items that would contain "soft" information. For instance if accurate information was to be obtained on smoking it should include the type of cigarette, the amount of nicotine contained in this brand, the number of cigarettes smoked, the gestation during which smoking continued and when the patient stopped smoking during pregnancy, if this was the case. Similar examples can be found with respect to alcohol and drug use, etc.

4. Outcome data

The interpretation of information from the prenatal sheet in

many instances requires knowledge of the final outcome of the pregnancy. This raises a number of difficulties with extracting material from hospital records, and again raises the question of accuracy with respect to Apgar scores and other measurements of outcome.

5. Analysis of data

The collection of information certainly implies its analysis and this raises the further question of which central agency would be the coordinator for the retrieval of this information; where would it be centralized and who would undertake any analysis?

For example, a nutritional history would certainly require more space and more specific questions than are currently present on any of the prenatal forms. The concept of data collection along these lines is more suggestive of a survey and would require more time than is realistically available to the practising physician.

It is extremely important, therefore, to clearly define the function of a prenatal record. Because a working document for the practising physician is incompatible with a document designed for accurate data collection, attempts at compromise to achieve both goals will almost certainly result in a document that does not satisfy either objective.

5. Summary and Conclusions

A review of the prenatal sheets in use across Canada shows that a common purpose is to function as a clinical document for the practising physician. Despite their apparent diversity all forms are fairly uniform in their content of basic obstetrical information and material. The inclusion of aids to assessing obstetric risk appears to have been accepted as a principle by almost all of the

provinces but there is very little uniformity with respect to the risk assessment systems in use. There is a wide diversity in design of most prenatal forms.

It would appear that general agreement could be reached upon the functions of a prenatal record and the inclusion of the necessary basic information that should be contained in a prenatal form as a minimum data base for patient management. The question of design will likely be much more difficult to resolve, particularly within the time constraints pending a final report of the Committee. The issue of using prenatal forms for data collection of a substantial nature is probably unrealistic although some agreement might be reached upon specific items of information, common to all forms, that might be collected on a provincial basis for comparison. The issue of risk assessment and the appropriate markers for such assessment is being reviewed by another sub-committee and the results of these deliberations could be incorporated into prenatal records if it was felt to be appropriate.

APPENDIX 4

LITERATURE REVIEW: FETAL HEART RATE MONITORING

APPENDIX 4

LITERATURE REVIEW: FETAL HEART RATE MONITORING

INTRA-PARTUM FETAL MONITORING

INTRODUCTION

Fetal heart tones have been accepted by clinicians as being important since the early 1800s. The use of auscultation and the recognition of fetal heart rate patterns were propounded by Kennedy in Dublin in a monograph published in 1833. In the latter half of the 19th century, Schwartz recommended that FHR be recorded during labour, both between and during contractions and suggested that fetal "asphyxia intoxication" modified fetal heart rate as did uterine contractions. Likewise, other physicians noted erratic patterns and the effect of drugs on these patterns about the same time. It is surprising, therefore, that these concepts largely disappeared until the 1950s when obstetricians again became interested in fetal heart patterns.

The few studies in the 1950s, using auscultation, suggested that the early definitions of fetal distress using fetal heart patterns were inadequate and poor predictors.

The use of electronic monitors has varied over previous decades, but recently has returned to prominence with improved equipment and the addition of fetal scalp sampling and other monitored variables. The prevailing philosophy in the 1980s appears to be that if bigger, better or more monitoring is done, it will be possible to identify all fetal distress and intervene in adequate time. How true is this?

Although several papers appeared in the late 1960s, it was not until the 1970s that real attention was paid to the evaluation of the usefulness of Fetal Heart Rate Patterns in the prediction of fetal distress. For this reason the literature has been reviewed from 1970 to the present day.

FORMAT OF LITERATURE REVIEW

Based on the same criteria used in the Task Force on the Periodic Health Examination, we looked at:

1. The burden of suffering
2. The validity and acceptability of manoeuvre
3. The effectiveness of the resulting intervention.

Number 2 has been divided further into the question posed in the original preliminary summary on fetal monitoring presented to this Committee in June. These are:

1. What is the objective of the risk indicator?
2. How well does it meet the objectives?
3. Who identifies the marker?
4. What are the costs?
5. What are the benefits?
6. What psychological cost is there?

The papers have been reviewed in reverse chronological order and have been divided into categories according to the strength of the research design and the quality of the evidence.

Burden of Suffering

The most common outcome measure suggested was perinatal mortality. In 1978, in Ontario, this accounted for 14.6/1000 deliveries. Although this appears high compared to other industrial nations, it is not common enough to be useful in any single series of cases. Because of the small denominators, any estimate of perinatal mortality is necessarily precarious and, without a confidence interval, can have little meaning on its own.

The second most frequent outcome suggested was Apgar Score which is used as a substitute measure of 'fetal distress'. It is

difficult to obtain the true incidence of low Apgar Scores (<7) in a total population since none of the authors carried out a community survey and hospital audits reflect the referral bias. The average estimate seems to be about 15% but this is probably high due to these biases.

Question 1: *WHAT IS THE OBJECTIVE OF THE INDICATOR?*

For fetal heart rate monitoring, the objective is to speedily identify the fetus in *distress* so that steps can be taken to prevent intra-partum death or the birth of an asphyxiated fetus, destined to have problems particularly in the first week of life. The implied philosophy is that a *distressed* fetus will be at high risk of perinatal death or a neurologically poor outcome resulting in handicap.

(The word 'distress' appears in italics because there is frequent argument concerning the difference between 'stress' and 'distress' and those two terms should be distinguished.)

Question 2: *HOW WELL DOES THE INDICATOR REACH ITS OBJECTIVE?*

The bulk of the papers deal with this question. There are two aspects to this question. The first (a) is how good is the indicator at doing its task and for this it will be necessary to look at fetal monitoring as a diagnostic test and explore its properties, sensitivity, specificity and predictive values. The second aspect (b) is to judge whether, despite its strengths and weaknesses, fetal monitoring as a diagnostic test has proved useful enough to alter the outcomes of delivery.

Under ideal test conditions, there should be 100% accuracy and no false positives or negatives, when gauged against a gold standard.

The gold standard used in the majority of papers was a low Apgar Score (< 7) and occasionally perinatal death. Sometimes, Apgar Score was associated with scalp pH.

Randomized Clinical Trials

Because of the use of comparable groups, the four RCT's did not address the problem of accuracy but went directly to the 2nd aspect of outcomes.

Cohort

The cohort studies likewise looked at comparability of outcomes and not at test itself.

Descriptive (Case Series)

The following table summarizes the papers in chronologically reverse order:

<u>1' Apgar <7</u>					
<u>Review</u>	<u>n</u>	<u>+Predictive Value</u>	<u>-Predictive Value</u>	<u>Sensitivity</u>	<u>Specificity</u>
1. Westgren et al.	5037	NC	NC	NC	NC
2. Phillips et al.	37	"	"	"	"
3. Liu et al.	50	"	"	"	"
4. Goodlin et al.	1100	"	"	"	"
5. Nochimson et al.	?	"	"	"	"
6. Amato	3076	"	"	"	"
7. Lee et al.	?	"	"	"	"
8. Saldana	620	23.0	86.2	70.6	43.7
9. Chik et al.	61	22.6	79.3	87.5	47.8
10. Tejani et al.	200	27.4	91.4	74.3	58.2
11. Tutela et al.	608	NC	NC	NC	NC
12. Edington et al.	911	"	"	"	"
13. Shenker et al.	760	39.0	77.8	53.4	66.2
	413	16.4	90.4	38.3	67.5
	1566	15.4	90.2	44.8	74.7
14. Thomas	946	NC	NC	NC	NC
15. Sturbois		68.4	91.4	76.5	87.6
	131	51.3	84.8	58.8	80.4
		52.8	84.2	55.9	82.5
16. Gabert et al.	396	NC	NC	NC	NC
17. Kelly et al.	150	"	"	"	"
18. Gabert et al.	749	66.2	91.4	84.3	79.5
19. Shenker	1000	24.5	89.7	68.4	56.6
20. Schiffrin et al.	307	42.9	93.0	53.9	89.6
21. Paul	1798	NC	NC	NC	NC
22. Paul et al.	245	"	"	"	"

Why is there such a difference in predictions? The main contribution to the variable predictive rates is the bias of referral. It is impossible to find details of the population except that, for the most part, the papers are dealing with high risk patients by their own criteria. The one exception is that of Westgren (#1) which looks at low risk and Amato (#6) who looks at a community hospital.

The worrisome aspect of these results is that, with one or two exceptions only, most of the authors were optimistic and enthusiastic proponents of monitoring despite positive predictive values dropping as low as 15%. The logical extension of these results is even more worrisome. Since the papers with low predictive values all studied high risk patient populations with a high prevalence of poor outcomes, things can only get worse if a 'normal' population is monitored since a lower prevalence will lead to more false positives being identified.

On the other hand, the high negative predictive values which were universally achieved are reassuring. This indicates that if the FHR tracings are normal then one can state with confidence that a normal fetus will result.

A summary of the methodological guidelines for a diagnostic test and the papers' approximation to those may provide a better appreciation of the problems. (See table on page 96).

Biases: Gold Standard

Apgar as a gold standard (best available estimate at the moment) may not be the best measure of fetal distress. Perinatal mortality is susceptible to small sample estimates.

Patient Sample Includes Spectrum of Disease

Most of the populations were 'high risk' (according to a variety of criteria) or low risk but only one is a community based hospital. As a result, the features of the test cannot be generalized.

Diagnostic Test

RCTs	Gold Standard	Patient Sample	Study Setting/Referral	Precision/Variation	Normal Defined	Technique Described	Utility of Test	Properties Discussed	Percentage Monitored
Haverkamp	X		X	X	X	X			
Kelso	X		X		X	X			
Renon	X		X						
Haverkamp	X		X		X	X			
<u>COHORT</u>									
Krebs	X	X			X	X	X	X	
Paul	X		X						
<u>DESCRIPTIVE</u>									
Westgren	X		X						
Phillips	X				X				
Liu	X				X				
Goodlin	X		X		X				
Nochimson	X				X	X			
Amato	X	X							
Lee	X		X		X				
Chik	X				X	X			
Tejani	X					X			
Tutela	X					X			
Edington	X		X		X				
Shenker	X				X	X			
Thomas	X				X				
Sturbois	X				X	X	X	X	
Gabert	X								
Kelly	X								
Gabert	X	X				X	X		
Shenker	X								
Shifrin	X				X	X	X	X	
Paul	X					X			

Study Sample/Referral Pattern

Not only does the referral pattern affect the outcome, it also contributes to an 'apparently' more accurate test if only high risk patients are included or a poor test if only low risk is considered. The second point is the percentage of total patients included. In many papers it is a small, perhaps carefully selected group of patients.

Precision/Variability

Only one paper discusses observer agreement and that is the clinical trial of Haverkamp. In any large series, if there exists intra- and inter-observer variability, it could under- or overestimate the value of the test.

Normal Defined

If normal/abnormal is not defined, it is impossible to assess the usefulness of the results.

Technique of Test

The same comment can be made for describing the test. Some of the papers carried a lot of detail which was helpful, others just 'monitored'.

Utility of Test

This was poorly documented. The bias resulting from its absence can be in the direction of overemphasizing the usefulness and raising false hopes in the interpreter of the results.

Properties

It was surprising that the obvious proponents of monitoring were blithely unaware of what their results were showing. Most did not discuss the problems of false positives and false negatives or discussed these as irrelevant. The bias of interpretation is strong.

Question 3: *WHO IDENTIFIES THE MARKER?*

The majority of papers make the point that personnel are trained both for the hooking up of the monitor and also the reading of the tracings. One paper goes as far as to point out that their unit has a fetal intensive care unit where a 'monitoring team' does all the intra-partum monitoring and this they consider essential.

How well are the tracings read? Haverkamp's second trial reports a 90% agreement between observer, but this is not borne out in a paper devoted to assessing tracings, written by Trimbos and Kierse. In their study, 100 tracings were presented twice to five readers in a random fashion and five components were assessed: Basal heart rate; oscillation amplitude and frequency; accelerations and decelerations and a sixth pooled index where 0, 1, 2 were assigned to each of the five components giving scores ranging from zero to 10. The following results were obtained:

Agreement	Basal HR		Oscillating Amplitude		Oscillating Frequency		Accel.		Decel.		Fischer's Score	
All 5	96	96	40	37	47	47	34	69	65	71	47	64
1 and 4	1	2	35	42	38	20	29	22	16	15	25	25
2 and 3	2	2	24	20	12	16	25	7	13	8	27	9
3,1,1, or 2,2,1	-	-	1	1	3	-	12	2	6	6	1	2

The accelerations were so variable that it had to be redefined in terms of amplitude and duration. Seven to 13% of the results were thought to warrant immediate intervention and then changed to normal on the second occasion.

The paper concludes that the interpretation of cardiotocogram (CTGS) is not as easy or reproducible as one might wish.

Question 4: *WHAT ARE THE COSTS?*

Banta and Thatcher, in their paper on the assessment of costs and benefits of electronic fetal monitoring, have estimated both direct and indirect costs and both are stated to be conservative. The direct cost of fetal monitoring looks at equipment services; neonatal and maternal morbidity due to monitoring as one component and direct and indirect cost of Cesarean section. It is difficult to judge how appropriate the costs are but they are estimated to run around \$411 million per year in the U.S.

Other papers have not tried to estimate monetary cost but have tabulated the amount of equipment which would be required for 100% monitoring. The estimates of a monitor to patient rates range from 1/300 - 1/500 patients.

The medical costs are better documented. The four randomized clinical trials reported increased c/s rates, with those of Haverkamp putting the rate as double that of an auscultated group.

The descriptive papers show variable Cesarean rates. Most suggest that it stays about the same, others claim a reduction and one or two show an increase. Unfortunately, without proper controls, it is hard to estimate.

Other medical costs to the mother can include lacerations of the mother or placenta and uterine perforation although these are rarely reported. Fetal costs include scalp lacerations; hemorrhage occurring in 0.3% cases and scalp abscesses. If amniotomy is done, a prolapsed cord is possible. More severe sequelae can occur, but are rare.

Cesarean sections can have risks for the fetus lung, but estimates of the extent of this are scanty.

Question 5: *WHAT ARE THE BENEFITS?*

The best evidence comes from the clinical trials because the patients had an equal chance of being auscultated or monitored. Of the four trials, three reported no benefit, but the fourth trial was stopped because it was felt that monitoring had proved to be superior. Unfortunately, the fourth trial had done frequent analysis of the data without an '*a priori*' set of stopping rules.

The cohort studies, like the descriptive studies reported an improvement in neonatal and perinatal mortality. Most also reported similar or fewer low Apgar scores. Lack of truly comparable groups make these claims impossible to assess.

Question 6: *WHAT ARE THE PSYCHOLOGICAL COSTS TO MOTHER?*

Some women reported anxiety and discomfort with the insertion of the electrodes, but often found the monitoring reassuring. One paper, by Starkman, enumerates the feelings of 35 women who had been monitored. Those who had had a previous fetal loss were positive about the benefits; those of low risk and no previous loss were predominantly negative.

SUMMARY

What conclusions can we come to in the light of the conflicting evidence? What follows is a series of quotes from the papers which have been evaluated, followed by a list of the recommendations from the NICHD Consensus Development Task Force Report: Predictors of Intrapartum Fetal Distress - The Role of Electronic Fetal Monitoring.

Quotes:

Paul (1974): 'To evaluate fetal monitoring fully, a well designed obstetric study combined with comprehensive infant growth and development studies is necessary.'

Kelso (1976): 'No beneficial or deleterious effects of continuous Fetal Heart Rate shown.'

Renou (1976): 'Highly trained staff necessary.'

Westgren (1980): 'Fetal Heart Rate Pattern irreproachable in less than 50% of tracings. Can mislead clinicians into unnecessary procedures.' (He advocates FHR monitoring but sees it as a screening device rather than a diagnostic test.)

Goodlin (1977): 'Prolapsed cord and abruptio placenta diagnosed not by FHR but by nursing care.'

Amato (1977): 'Although differences seen, they are usually in the lower birthweights. Once over 2500 grams, the differences disappear.' (This may be one explanation why the clinical trials of Haverkamp showed no differences - he did not monitor less than 34 weeks GAs.)

Lee (1976): 'Fetal monitoring can predict a good outcome but seems to overestimate problems.'

Tutera (1975): 'Technical difficulties and complications can occur.'

Edington (1975): '20% of bad outcomes cannot be predicted by FHR.' (This point also made by Paul in one of his papers where he explored the physiological and biological reasons why intrapartum monitoring could fail - uterine contractions unpredictable, cord position, etc.)

Shenker (1975): 'Statistically, the method of delivery is almost as accurate a predictor of low APGAR scores at one minute as monitoring record.'

Thomas (1975): 'Late decelerations are frequently temporary and not serious.' Monitoring is good screening, not a diagnostic device.

NICHD Consensus Development Task Force

Recommendations for current clinical practice:

1. The electronic fetal monitor or any other technology should never be a substitute for clinical judgment. Electronic fetal monitoring is only one method of fetal assessment.
2. Proper use of both intermittent auscultation and continuous electronic fetal monitoring in both high- and low-risk patients should, at the outset, include a discussion with the patient of her wishes, concerns and questions concerning benefits, limitations, and risks of fetal monitoring. Women should have the opportunity to discuss the use of all forms of monitoring during the course of prenatal care, and again upon admission to the labour suite.
3. Periodic auscultation of the fetal heart rate is an acceptable method of assessment of fetal condition for women at low risk of intrapartum fetal distress. Interpretation of auscultated fetal heart rate data should include an understanding of the relationship of FHR changes to uterine contractions. Although the Task Force finds no evidence that electronic fetal monitoring reduces morbidity and mortality in low-risk patients, it recognizes that, under certain circumstances, mothers or physicians may choose to use electronic fetal monitoring even in low-risk situations.
4. The use of electronic fetal monitoring should be strongly considered in high-risk patients. Some of the high risk situations may include: (a) low birth weight, prematurity, postmaturity, and IUGR; (b) medical complications of pregnancy; (c) meconium staining of the amniotic fluid; (d) intrapartum obstetrical complications; (e) use of oxytocin in labour; and (f) the presence of abnormal auscultatory findings.

The medical record should reflect careful consideration of the benefits and risks to each patient, including a discussion of the indications for electronic fetal monitoring with the patient.

5. Since unexpected risk factors may arise during labour in patients without prior evidence of risk, all hospitals and birthing centres providing maternity care should have the necessary trained staff and equipment to assess carefully the status of each fetus in labour and to take appropriate action.
6. In order that electronic fetal monitoring be used appropriately, the medical profession and others should encourage, through their various educational modalities a thorough understanding of the principles and procedures of intrapartum FHR assessment by personnel responsible for the care of pregnant women. Special attention should be given to benefits, limitations and risks.
7. The use of fetal scalp blood pH determination is strongly encouraged as an adjunct to electronic FHR monitoring.
8. Attention to the known potential hazards of electronic fetal monitoring should accompany its use. Placement of the fetal scalp electrode and intrauterine pressure catheter should be performed with attention to aseptic and atraumatic technique. Prolonged supine position of the mother should be avoided, and maternal mobility should not be unnecessarily limited.
9. Hospital personnel should be cognizant of the potential impact of electronic fetal monitoring upon family centered childbirth. Family-centered care and indicated intrapartum electronic fetal monitoring are not mutually exclusive. Maternity services should be encouraged to integrate concepts of family-centered care with care of women who are electronically monitored.

FETAL HEART RATE

RANDOMIZED CLINICAL TRIALS

Langendoerfer, S.; Haverkamp, A.D.; Murphy

"Pediatrics follow-up of a randomized controlled trial of intrapartum fetal monitoring techniques". Journal of Pediatrics 1980 97: 1, 103-107.

Haverkamp, A.D.; Orleans, M.; et al.

"A controlled Trial of the Differential Effects of Intrapartum Fetal Monitoring". American Journal of Obstetrics and Gynecology 1979 134: 399.

Kelso, I.; Parsons, J.

"An assessment of continuous Fetal Heart Rate monitoring in labour: A randomized clinical trial". American Journal of Obstetrics and Gynecology 1976 131: 5, 526-532.

Trimbos, J.B.; Keirse, M.J.

"Assessment of Ante-partum Cardio-Tocograms in High Risk Pregnancy". British Journal Obstetrics & Gynecology, 1980, April: 87(4): 261-269.

Renou, P.; Chang, J.; Anderson, I.; Wood, C.

"Controlled trial of fetal intensive care". American Journal of Obstetrics and Gynecology 1976 126: 4, 470.

Haverkamp, A.D.; Thompson, H.E.; McFee, J.G.; Cetrulo, C.

"The evaluation of continuous fetal heart rate monitoring in high risk pregnancy". American Journal of Obstetrics and Gynecology 1976 125: 310.

COHORT

Keane, M.W.D.; Horger, E.O.; Vice, L.

"Comparative study of stressed and non-stressed antepartum fetal heart rate testing". Obstetrics and Gynecology 1981 March 57(3): 320-4.

Krebs, H.B.; Petres, R.E.; Dunn, L.J.; Segretti, A.

"Intrapartum fetal heart monitoring". American Journal of Obstetrics and Gynecology 1980 September 138(2): 213-9.

Paul, R.H.; Hon, E.H.

"Clinical fetal monitoring. V. Effect on perinatal outcome".
American Journal of Obstetrics and Gynecology 1974 February
118(4): 529-33.

CASE SERIES

Westgren, M.; Ingemarsson, E.; Ingemarsson, I.; Solum, T.

"Intrapartum Electronic Fetal Monitoring in Low Risk Patients"
Obstetrics and Gynecology 1980, September; 56(3): 301-304.

Phillips, W.D.; Towell, M.E.

"Abnormal Fetal Heart Rate Associated with Congenital Abnormalities"
British Journal of Obstetrics and Gynecology 1980, April; 87(4)
270-274.

Liu, D.T.Y.; Blackwell, R.J.; Tukel, S.

"The Relevance of (antenatal) and Intrapartum Fetal Heart Rate
Patterns to Fetal Outcomes."
British Journal of Obstetrics and Gynecology 1978; 85: 270-277

Goodlin, R.C.; Haesslin, H.C.

"When is it Fetal Distress?" American Journal of Obstetrics and
Gynecology 1977; 128(4) 440-447.

Nochimson, D.J.; Cetrulo, C.L.

"Intrapartum Fetal Monitoring" Journal of Family Practice 1977;
4(5): 845.

Amato, J.C.

"Fetal Monitoring in a Community Hospital" Obstetrics and
Gynecology 1977; 50: 269-274.

Lee, W. .; Baggish, M.S.

"The Effect of Unselected Intrapartum Fetal Monitoring" Obstetrics and Gynecology 1976; 37(5): 516-520.

Chik, L.; Sokol, R.J.; Rosen, M.G.

"Prediction of the One Minute APGAR Score from Fetal Heart Rate Data" Obstetrics and Gynecology 1976; 48(4): 452-455.

Tejani, N.; Mann, L.I.

"Correlation of Fetal Heart Rate Patterns and Fetal pH with Neonatal Outcome" Obstetrics and Gynecology 1976; 48(4): 460-63.

Tutera, G.; Newman, R.L.

"Fetal Monitoring: Its Effect on Perinatal Mortality and Cesarean Section and its Complications" American Journal of Obstetrics and Gynecology 1975; 122: 750-54.

Edington, P.T.; Sibanda, J.; Beard, R.W.

"Influence on Clinical Practice of Routine Intrapartum Fetal Monitoring" British Medical Journal 1975; 3: 341-3.

Shenker, L.; Post, R.C.; Seiler, J.S.

"Routine Electronic Monitoring of Fetal Heart Rate and Uterine Activity During Labour" Obstetrics and Gynecology 1975; 46(2): 185-89.

Thomas, G.

"The Aetiology, Characteristics, and Diagnostic Relevance of Late Deceleration Patterns in Routine Obstetric Practice." British Journal of Obstetrics and Gynecology 1975; 82: 121.

Sturbois, G.; Tournaire, M.; Breart, G.

"Prediction of one minute APGAR Score by Fetal Heart Rate Analysis: A Comparison Between Three Methods" International Journal of Gynecology and Obstetrics 1975; 13: 118-122.

Gabert, H.A.; Stenchever, M.A.

"Electronic Fetal Monitoring as a Routine Practice in an Obstetrical Service: A Progress Report" American Journal of Obstetrics and Gynecology 1974, 118(4): 534-7.

Kelly, V.C.; Kulkarni, D.

"Experiences with Fetal Monitoring in a Community Hospital"
Obstetrics and Gynecology 1973; 41(6): 818-824.

Gabert, H.A.; Stenchever, M.A.

"Continuous Electronic Monitoring of Fetal Heart Rate during Labour"
American Journal of Obstetrics and Gynecology 1973; 115(7): 919-923.

Shenker, L.

"Clinical Experiences with Fetal Heart Rate Monitoring of 1000 Patients in Labour" American Journal of Obstetrics and Gynecology 1973; 115(8): 1111-6.

Schiffrin, B.S.; Dame, L.

"Fetal Heart Rate Pattern" Journal of American Medical Association 1972; 219(10): 1322-5.

Paul, R.H.

"Clinical Fetal Monitoring" American Journal of Obstetrics and Gynecology 1972; 113(5): 573-77.

Paul, R.H.; Hon, E.H.

"A Clinical Fetal Monitor" Obstetrics and Gynecology 1970; 35(2): 161-9.

REVIEWS AND EDITORIALS

Dixon, A.S.

"The Electronic Fetal Monitor: Should Every Mother Have one?"
Canadian Family Physician 1981, June; 27: 1023-28.

Hill, L.M.

"Diagnosis and Management of Fetal Distress" Mayo Clinic
Proceedings 1979, December; 54 (12): 784-93.

Hobbins, J.C.; Freeman, R; Queenan, J.T.

"The Fetal Monitoring Debate" Obstetrics and Gynecology 1979,
July; 54(1): 103-109.

Goodlin, R.C.

"History of Fetal Monitoring" American Journal of Obstetrics
and Gynecology 1979, February; 133 (3): 323-52.

Banta, H.D.; Thacker, S.B.

"Assessing the Costs and Benefits of Electronic Fetal Monitoring"
Obstetrics and Gynecology Survey, 1979, 34 (8): 627-42.

Editorial

"Intrapartum Fetal Monitoring For all?" British Medical Journal,
1976; II: 1466.

Editorial

Obstetrical Gynecological Survey, 1975; 30: 503-504.

Zuspan, F.P.; Quilligan, E.J.; Iams, J.D.; Van Geijn, H.P.

"NICHD Consensus Development Task Force Report: Predictors of
Intrapartum Fetal Distress - The Role of Electronic Fetal
Monitoring" Editorial, American Journal of Obstetrics and Gynecology
1979, October; 135 (3): 287-91.

COST BENEFIT

Banta, H.D.; Thacker, S.B.

"Assessing the Costs and Benefits of Electronic Fetal Monitoring"
Obstetrical and Gynecological Survey 1979; 34 (8): 627-42.

Neutra, R.R.; Fienberg, S.E., et al.

"Effect of Fetal Monitoring on Neonatal Death Rates"

New England Journal of Medicine 1978; 299: 324-26.

Quilligan, E.J.; Paul, R.H.

"Fetal Monitoring - Is it Worth it?" Obstetrics and Gynecology

1975, January; 45 (1): 96-100.

PSYCHOLOGICAL COSTS

Starkman, M.N.

"Fetal Monitoring: Psychological Consequences and Management
Recommendations" Obstetrics and Gynecology 1977; 50: 500-504.

Youngs, D.D.; Starkman, M.N.

"Psychological and Physiological Aspects of Electronic Fetal
Monitoring" Primary Care 1976; 3: 691-700.

APPENDIX 5

PROVINCIAL PRENATAL RECORD FORMS

PART I

HOSPITAL

BRITISH COLUMBIA

1. NAME				RACIAL ORIGIN		BRITISH COLUMBIA			
ADDRESS				TELEPHONE NUMBER		DOCTOR'S NAME			
MOTHER'S AGE		DATE OF BIRTH		OCCUPATION				MARITAL STATUS	
FATHER'S NAME				RACIAL ORIGIN					
FATHER'S AGE		FATHER'S OCCUPATION		M S P NUMBER		DOCTOR'S PHONE NUMBER			

2. OBSTETRICAL HISTORY INCLUDING ABORTIONS

[illegible]

3. RISK ASSESSMENT

Reproductive History

- ☐ Age - less than 15, greater than 35
- ☐ Parity 5 +
- ☐ Weight - less than 45, greater than 90kg (< 100, > 200 lbs)
- ☐ Habitual Abortion
- ☐ Stillbirth/Neonatal Death
- ☐ Fetal Anomalies
- ☐ Labour Premature
- ☐ Labour Prolonged
- ☐ Labour precipitous (< 3 hours)
- ☐ Difficult Delivery
- ☐ Post p. Hemorrhage, Manual Removal
- ☐ Preeclampsia/Hypertension

Associated Hi Risk Conditions

- ☐ Uterine Surgery or Malformations
C-section, Cone Biopsy,
Cervical Suture, etc.
- ☐ Chronic Renal Disease
- ☐ Diabetes (incl. Gest.)
- ☐ Cardiac Disease & Hypertension
- ☐ Serious Medical, Surgical or
Psychiatric Disorders

4. HISTORY OF PRESENT PREGNANCY (Specify)

NO	YES
☐ BLEEDING (Date)	☐ _____
☐ PYREXIA (Date)	☐ _____
☐ RADIATION (Date)	☐ _____
☐ NAUSEA	☐ _____
☐ SMOKING	☐ _____
☐ (Query pattern of usage)	☐ _____
☐ ALCOHOL DRUGS (Query pattern of usage)	☐ _____
☐ OCCUPATIONAL	☐ _____
NUTRITION	
(24 hour recall) GOOD ☐	FAIR ☐ POOR ☐

5. PAST ILLNESS (Specify)

NO	YES
☐ RENAL	☐
☐ CARDIAC	☐
☐ INFECTIONS	☐
☐ RUBELLA	☐
☐ VENEREAL	☐
☐ ALLERGIES	☐
☐ OPERATIONS	☐
☐ TRANSFUSIONS	☐
☐ OTHERS	☐

6. FAMILY HISTORY (Specify)

NO	YES
☐ DIABETES	☐
☐ CARDIAC	☐
☐ HYPERTENSION	☐
☐ TUBERCULOSIS	☐
☐ TWINS	☐
☐ MALFORMATIONS	☐
☐ OTHERS	☐

10. EXAMINATION

DESCRIPTION	CONDITION
HEAD & NECK	
BREAST & LYMPH NODES	 mm Hg
HEART & LUNGS	
ABDOMEN	
SPINE	
VARICES & SKIN	
PELVIS NORMAL	
ABNORMAL (Describe)	
CERVIX CYTOLOGY (Class No.)	
OTHER OBSERVATIONS	

7. MENSTRUAL HISTORY

LMP (Date)
EDC
IF UNSURE, CONSIDER EARLY ULTRASOUND
MENSTRUAL CYCLE

8. METHOD OF CONTRACEPTION
(Specify)

DATE OF DISCONTINUANCE	
------------------------	--

9. CURRENT MEDICATIONS

[illegible]

11. TOPICS FOR DISCUSSION AND ADVICE

Alternate Physician(s)	Bowels
Genetic Counseling	Baths
Prenatal Course	Clothes
& Classes	Books
Nutrition	Breast Care
Supplements	Breast Feeding
Wt. Control	Baby Care
Exercise	Labour Stages
Hosp. Adm. Procedures	Rest
Arrival at Hospital	Smoking
Anaes. & Analg.	Alcohol
Coitus	Medications

LABOUR SUMMARY AND DELIVERY RECORD PART 1

1	GRAV	PARA	ABORTIONS SPONTANEOUS ☐ ELECTIVE ☐		
	LMP	CYCLE	QUICKENING		
	EDC		SURE ☐ UNSURE ☐		GESTATION
	BLOOD GP	LAST TITRE	DATE	LAST Hgb	DATE
					RUBELLA TITRE

2 PREGNANCY SINGLE ☐ TWINS ☐ TRIPLETS ☐ OTHER ☐ SPECIFY _____	In Multiple Pregnancy specify baby A, B, C, etc. (Use separate form for each baby)
--	---

3	ADMISSION TO LABOUR/ Time _____ In True Labour ☐ In Suspected Labour ☐ For Induction ☐ For Elective C. Section ☐ DELIVERY AREA Date _____ Other ☐ Specify _____
---	---

4	HIGH RISK FACTORS BEFORE LABOUR SCORES (if applicable) - Present Pregnancy <table border="1" style="width: 40px; height: 40px; display: inline-table;"></table> - Obstetrical History <table border="1" style="width: 40px; height: 40px; display: inline-table;"></table> - Gestational Age <table border="1" style="width: 40px; height: 40px; display: inline-table;"></table> TOTAL SCORE <table border="1" style="width: 40px; height: 40px; display: inline-table;"></table>	LIST:-- _____ _____ _____ _____
---	---	--

5	CONSULTATION OBSTETRIC <table border="1" style="width: 100px; height: 40px; display: inline-table;"> <tr><td>NONE BEFORE LABOUR</td></tr> <tr><td>DURING LABOUR</td></tr> <tr><td>AFTER LABOUR</td></tr> </table> TIME CALLED (Hour) <table border="1" style="width: 60px; height: 20px; display: inline-table;"></table>	NONE BEFORE LABOUR	DURING LABOUR	AFTER LABOUR	PEDIATRIC <table border="1" style="width: 100px; height: 40px; display: inline-table;"> <tr><td>NONE BEFORE LABOUR</td></tr> <tr><td>DURING LABOUR</td></tr> <tr><td>AT DELIVERY</td></tr> </table> TIME CALLED (Hour) <table border="1" style="width: 60px; height: 20px; display: inline-table;"></table>	NONE BEFORE LABOUR	DURING LABOUR	AT DELIVERY
NONE BEFORE LABOUR								
DURING LABOUR								
AFTER LABOUR								
NONE BEFORE LABOUR								
DURING LABOUR								
AT DELIVERY								

6	INDUCTION OR AUGMENTATION OF LABOUR INDUCTION No ☐ Yes ☐ AUGMENTATION No ☐ Yes ☐ METHOD OF INDUCTION OR AUGMENTATION ARM Only ☐ Oxytocin Only ☐ ARM and Oxytocin ☐ RIPENING INFUSIONS No ☐ Yes ☐ Other ☐ Specify _____ INDICATIONS FOR INDUCTION ☐ _____ OR AUGMENTATION ☐ _____
---	--

7	FETAL MONITORING - FETAL HEART Auscultation or other means On Admission <table border="1" style="width: 100px; height: 20px; display: inline-table;"></table> 1st Stage <table border="1" style="width: 100px; height: 20px; display: inline-table;"></table> 2nd Stage <table border="1" style="width: 100px; height: 20px; display: inline-table;"></table>	FETAL BLOOD FBS No ☐ Yes ☐ No. of Times <table border="1" style="width: 40px; height: 20px; display: inline-table;"></table> Lowest pH <table border="1" style="width: 40px; height: 20px; display: inline-table;"></table> Cord Blood Gases No ☐ Yes ☐ If Yes, UApH <table border="1" style="width: 40px; height: 20px; display: inline-table;"></table>	ELECTRONIC MONITORING (Excluding Doppler) Continuous ☐ <table border="1" style="width: 100px; height: 20px; display: inline-table;"></table> Intermittent ☐ FHR <table border="1" style="width: 100px; height: 20px; display: inline-table;"></table> TIME FROM <table border="1" style="width: 60px; height: 20px; display: inline-table;"></table> TO <table border="1" style="width: 60px; height: 20px; display: inline-table;"></table>
---	---	---	---

8	TIME SUMMARY <table border="1" style="width: 100%; height: 100px; display: inline-table;"> <tr><th>Hour</th><th>Day</th><th>Month</th><th>Year</th></tr> <tr><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td></tr> </table>	Hour	Day	Month	Year																																					<table border="1" style="width: 100%; height: 100px; display: inline-table;"> <tr><th>Hours</th><th>Minutes</th></tr> <tr><td></td><td></td></tr> <tr><td></td><td></td></tr> <tr><td></td><td></td></tr> <tr><td></td><td></td></tr> <tr><td></td><td></td></tr> <tr><td></td><td></td></tr> <tr><td></td><td></td></tr> <tr><td></td><td></td></tr> <tr><td></td><td></td></tr> </table>	Hours	Minutes																		
Hour	Day	Month	Year																																																											
Hours	Minutes																																																													

9	BABY Unit Number <table border="1" style="width: 100px; height: 20px; display: inline-table;"></table> SEX - Male ☐ Female ☐ Ambiguous ☐ RECORD APGAR SCORE ON REVERSE OF WHITE COPY WEIGHT <table border="1" style="width: 60px; height: 20px; display: inline-table;"></table> GMS CONGENITAL MALFORMATIONS No ☐ Yes ☐ Specify _____	Admission No. <table border="1" style="width: 100px; height: 20px; display: inline-table;"></table> Liveborn ☐ Stillborn ☐ GESTATIONAL AGE <table border="1" style="width: 40px; height: 20px; display: inline-table;"></table> WEEKS Duration of Ruptured Membranes <table border="1" style="width: 100px; height: 20px; display: inline-table;"></table>
---	--	---

10	UMBILICAL CORD: PROLAPSE ☐ OCCULT ☐ TRUE KNOT ☐ LOOPS OF CORD No ☐ Yes ☐ VESSELS <table border="1" style="width: 40px; height: 20px; display: inline-table;"></table> <table border="1" style="width: 40px; height: 20px; display: inline-table;"></table> SPECIFY NUMBER AND SITE OF LOOPS _____ PLACENTA - Normal ☐ Abnormal ☐ Specify _____ WEIGHT <table border="1" style="width: 60px; height: 20px; display: inline-table;"></table> GMS TO PATHOLOGY ☐ CULTURES ☐ OTHER ☐ _____ CORD BLOOD - VD ☐ RH STUDIES ☐ RESERVE ☐ OTHER ☐ _____ DOCTORS PRESENT AT DELIVERY _____ NURSES PRESENT AT DELIVERY _____ SIGNATURE OF ATTENDING NURSE _____ DATE _____
----	---

WHITE COPY—MOTHER'S CHART YELLOW COPY—INFANT'S CHART PINK COPY—PHYSICIAN'S COPY

LABOUR SUMMARY AND DELIVERY RECORD

PART 2

11 INDICATIONS FOR OPERATIVE DELIVERY

(Enter in Section 15)

2nd Stage Delay
— Primigravida > 1 hour
— Multigravida > 1/2 hour
Fetal Distress
— Abnormal FHR
— Meconium
— Acidosis
Cord Prolapse or Presentation
Poor Maternal Effort
Arrest in 2nd Stage
— Transverse
— POP
Failed Trial of Forceps
Disproportion
Excessive Fetal Size
Prolonged Labour

Incoörd Uterine Action
Cervical Dystocia
Malpresentation
— Breech
— Face/Brow
— Shoulder, etc.
Unstable Lie
IUGR
APH
— Abruptio
— Plac Praevia
— Undiag Haem.
Cardiac Disease
Pre-Eclampsia/Eclampsia
Hypertension

Diabetes
Renal Disease
Immunization
— RH, etc.
Multiple Pregnancy
Elderly Primigravida
Congenital Malforms.
Known Fetal Anomaly
Prev. Uterine Surgery
— C Section (Repeat)
— Myomectomy
— Hysterotomy
Prev. Surgery of
Lower Genital Tract
Infections (Herpes,
Chorioamnionitis, etc.
Other

THIS RECORD FOR BABY

(A, B, C etc)

12 ANALGESIA DURING LABOUR

NARCOTIC No ☐ Yes ☐

Epidural Catheter ☐ Nitrous Oxide ☐

Other ☐ Specify _____

MEDICATION

IV/IM

DOSE

TIME

DATE

13 ANAESTHESIA DURING DELIVERY

Nitrous Oxide ☐

General Anaesthetic ☐

Spinal ☐

Epidural ☐

Pudendal ☐

Local Perineal ☐

Other ☐ Specify _____

14 HIGH RISK FACTORS DURING LABOUR AND DELIVERY

Meconium ☐
FHR Abnormal 1st Stage ☐
FHR Abnormal 2nd Stage ☐

Cord Complications ☐
Intrapartum Hypertension ☐
Intrapartum Eclampsia ☐
Intrapartum Haemorrhage ☐
Intrapartum Infection ☐

Multiple Pregnancy ☐
Malpresentation ☐
Prolonged ROM ☐
Prolonged Labour ☐
Feto Pelvic Disproportion ☐

Prolonged 2nd Stage ☐
Fetal pH < 7.25 ☐
UAPH < 7.20 ☐
Other ☐ _____
None ☐

15 DELIVERY

PRESENTATION AND POSITION — During Labour _____

At Delivery _____

Type	Spontaneous Vaginal ☐	Ventouse Extraction ☐	Shoulder Dystocia ☐
	Low ☐	Low Mid ☐	Mid ☐ High ☐
Forceps	Traction Mild to Mod ☐	Traction Mod to Severe ☐	
	Trial Successful ☐	Trial Failed ☐	
Rotation	Manual & Spontaneous Delivery ☐	Forceps Rotation & Spontaneous Delivery ☐	
	Manual & Forceps Delivery ☐	Forceps Rotation & Forceps Delivery ☐	
Breech	Easy ☐	Moderately Difficult ☐	Difficult ☐
	Internal Version ☐	Breech Extraction ☐	Forceps to after coming head ☐
Caesarean	Plannestiel Incision ☐	Vertical Midline Incision ☐	Paramedian Incision ☐ Tubal ligation ☐
Section	Low Segment Transverse Incision ☐	Low Segment Vertical Incision ☐	Classical ☐ Inverted-T ☐

OTHER TYPE OF DELIVERY ☐ Specify _____

MANUAL REMOVAL OF PLACENTA

No ☐ Yes ☐

INDICATIONS FOR OPERATIVE DELIVERY (To be completed for all Forceps, Rotations, Breech Extractions, Ventouse & Caesarean Sect's.)

List in Order of Priority 1. _____
2. _____
3. _____
4. _____

16 BLOOD LOSS

Minimal ☐

Moderate ☐

Excessive ☐

ESTIMATED

CC's

OR MEASURED

CC's

PERINEUM

Intact ☐

Tear

(DEGREE: Definitions on Reverse)

EPISIOTOMY

None ☐

Midline

Medio-lateral ☐

Extended into Vagina ☐

Extended in Rectum ☐

Additional Lacerations (Labial, Vaginal, Cervical, etc.)

No ☐

Yes ☐

If YES specify _____

Sutured By _____

Suture Pack left in No ☐ Yes ☐

Catherized No ☐ Yes ☐

Catheter left in: No ☐ Yes ☐

Additional Comments and Operative Findings: _____

SIGNATURE OF ATTENDING PHYSICIAN _____

DATE _____

NEWBORN RECORD - PART 1

1. HOSPITAL OF BIRTH:	
DOCTOR:	CASE NO.
S.O.C.	
FATHER'S NAME	RACIAL ORIGIN
MOTHER'S MAIDEN NAME	UNIT NUMBER
RACIAL ORIGIN	

2. FETAL RISK ASSESSMENT:	SCORES
NO PRENATAL RECORD ☐ PRENATAL RISK: None ☐ Yes ☐	
NO LABOUR/DELIVERY RECORD ☐ LABOUR/DELIVERY RISK: None ☐ Yes ☐	

3. MATERNAL HISTORY SUMMARY:		AGE	BLOOD GP.	RH POS.	RH NEG.
Number of previous pregnancies _____					
Term _____	Term _____	Therapeutic _____	Living _____		
LIVE BIRTHS: Pre-term _____	STILLBIRTHS: Pre-term _____	ABORTIONS: Spontaneous _____	NO. OF CHILDREN At Home _____		
COMMENTS: _____					

4. DELIVERY: DURATION MEMBRANE RUPTURE _____ Hrs. MECONIUM STAINING: No ☐ Yes ☐ Method of Delivery _____	
DATE OF BIRTH _____ Time _____ Hrs. SEX _____	CORD BLOOD SPECIMEN: VDS ☐ Rh ☐ Reserve ☐
COMMENTS: _____	

5. APGAR SCORE		6. RESUSCITATION:		7. EYE PROPHYLAXIS:	
	0	1	2	1 Min.	5 Min.
Heart Rate	Absent	Under 100	Above 100		
Resp. Effort	Absent	Slow Irregular	Good Crying		
Muscle Tone	Limp	Some flex of Extrem.	Active Motion		
Reflex Irritability	None	Grimace	Crying		
Colour	Blue Pale	Blue Extrem.	All Pink		
TOTAL					

6. RESUSCITATION:		7. EYE PROPHYLAXIS:	
No ☐ Yes ☐ By Whom: _____		A ₉ NO ₃ ☐ Other ☐	
Method: _____	Age (min) _____	Specify: _____	
☐ O ₂ Flow _____	Duration (min) _____	Signature: _____	
☐ O ₂ Mask & Pressure _____		8. INFANT IDENTIFIED BY: _____	
☐ O ₂ Endotracheal Tube _____		Comments: _____	
☐ Other _____			
Comments: _____			

9. DELIVERY ROOM ASSESSMENT:	
Colour: ☐ Pink ☐ Plethoric ☐ Central Cyanosis ☐ Pale Skin: ☐ Vernix ☐ Desquamation ☐ Meconium ☐ Edema ☐ Petechiae	
Activity: ☐ Active ☐ Lethargic Tone: ☐ Good ☐ Fair ☐ Limp/Floppy CVS: Apex Left ☐ Yes ☐ No	
Respiratory System: ☐ Normal ☐ Grunting ☐ Retraction ☐ Shallow Respirations ☐ Poor Air Entry ☐ Gasping	
Heart Sounds: ☐ Normal ☐ Abnormal Umbilical Cord: ☐ Normal ☐ Thick ☐ Thin ☐ Meconium Stained Vessels: ☐ 2 ☐ 3	
Abdomen: ☐ Normal ☐ Scaphoid ☐ Masses _____	
Genitalia: ☐ Female ☐ Male ☐ Ambiguous Anus: ☐ Patent ☐ Imperforate	
Birth Trauma: ☐ No ☐ Yes _____	
Congenital Anomalies: ☐ No ☐ Yes _____	
Immediately before Transfer to Nursery: IN O ₂ : ☐ No ☐ Yes Axillary Temp. _____ °C Resp. Rate _____ Heart Rate _____ Voided: ☐ Yes ☐ No	
Passed Meconium: ☐ Yes ☐ No EXAMINER'S SIGNATURE: _____	

10. TRANSFER TO:	
Transitional Nursery ☐ Regular Nursery ☐ Intensive Care Nursery ☐ Other ☐	
Time of Admission to Nursery: _____ Hours Date: _____	

11. EVALUATION OF DEVELOPMENT: (Growth Curves on Reverse)		THIS INFANT CLASSIFIED AS:	
Birthweight _____ (Gms) _____ % ILE	Length _____ (Cms) _____ % ILE	Small for Gestational Age	SGA ☐
Head Circ. _____ (Cms) _____ % ILE		Appropriate for Gestational Age	AGA ☐
		Large for Gestational Age	LGA ☐

GESTATION AGE:		Pre Term (< 37 Completed Weeks - i.e. < 259 Days) ☐	
By Dates _____ Wks. By Exam _____ Wks.	Method _____	Term (37 to < 42 Completed Weeks - 254 to 293 Days) ☐	
		Post Term (42 Completed Weeks or More - 294 Days or More) ☐	

WHITE COPY—INFANT'S CHART YELLOW COPY—MOTHER'S CHART PINK COPY—PHYSICIAN'S COPY

NEWBORN RECORD - PART 2

12. PROCEDURES:

SKIN CARE: No ☐ Yes ☐ _____
 CORD CARE: No ☐ Yes ☐ _____
 VIT. K₁: ☐

13. THIS INFANT TRANSFERRED TO ANOTHER HOSPITAL:

No ☐ Yes ☐ SPECIFY _____

14. PHYSICAL EXAMINATION: A. Admission B. Discharge

A. Date: _____ B. Date: _____ Axillary Temp. _____ °C Resp. Rate _____ /min. Apex Rate _____ /min.
 1 ☐ Appearance 7 ☐ Reflexes 13 ☐ Skin COMMENTS: (Use Ref. No's.)
 2 ☐ Abdomen 8 ☐ Eyes 14 ☐ Genitals
 3 ☐ Femoral Pulses 9 ☐ E.N.T. 15 ☐ Anus
 4 ☐ Heart 10 ☐ Head/Neck 16 ☐ Trunk/Spine
 5 ☐ Lungs 11 ☐ Extremities 17 ☐ Cry
 6 ☐ Tone 12 ☐ Hips 18 ☐ Cong. Anom.
 (CODE: ☒ = Normal ☐ = Abnormality Present (Describe) Signature: A. _____ /B. _____

15. PROBLEM LIST:

Date Commenced	PROBLEM	Resolved at Time of Discharge Yes/No/Unknown

16. ANY FURTHER PERTINENT INFORMATION:

17. POSSIBLE FOLLOW-UP PROBLEMS:

(CODE: ☐ = Problem Present (describe))
 COMMENTS: (Use Ref. No.) 
 A. Socio-Economic B. Emotional C. Infant
 1 ☐ Drug Habituation 8 ☐ Maternal Rejection 16 ☐ Unresolved Medical Problem
 2 ☐ Language Problem 9 ☐ Paternal Rejection 17 ☐ Potential Medical Problem
 3 ☐ Malnutrition Prolonged Maternal/ 18 ☐ Other Infant Problem
 Home Situation: 10 ☐ Infant Separation
 4 ☐ Single Parent 11 ☐ Maternal Immaturity D. Sibling with Chronic
 5 ☐ Working Mother 12 ☐ Maternal Anxiety State Medical Problem
 6 ☐ Poverty 13 ☐ Maternal Psychosis 19 ☐ "SIDS"
 7 ☐ Other Socio-Economic 14 ☐ Post Partum Depression 20 ☐ Other Chronic Medical
 15 ☐ Other Emotional Problem

18. FURTHER PROCEDURES: (See also Section 12)

METABOLIC SCREENING TESTS: No ☐ Yes ☐ Specify _____ CORD VDS TEST: Neg. ☐ Pos. ☐ URINALYSIS: Neg. ☐ Pos. ☐
 CIRCUMCISION: No ☐ Date: _____ Method: _____
 Yes ☐

19. CONSULTATIONS: With:

1. _____
 2. _____

20. NEONATAL DEATH: ☐

Date: _____ Time: _____ Hrs.

AUTOPSY: Planned ☐ Requested ☐
 Refused ☐

Comments: _____

21. CONDITION ON DISCHARGE:

DISCHARGED ALIVE ☐ Date: _____ Weight _____ GMS
 FINAL DIAGNOSIS: _____

DISCHARGED WITH: Mother ☐ Other ☐

Referral to Public Health Nurse: _____

Referral to Dept. of Human Resources: _____



Breast Fed ☐ OR Specify Formula: _____

Vitamins ☐ Fluoride ☐ Iron ☐ DISCHARGE MEDICATIONS: (Specify & state Dose) _____

22. DISCHARGED TO THE CARE OF:


ATTENDING PHYSICIAN: _____ M.D.

INTRAUTERINE GROWTH CHART



LUBCHENCO L.O. HANSMAN C. and BOYD E. Pediatrics 37:403 1966
 Copyright American Academy of Pediatrics 1966 and 1967

CLASSIFICATION OF NEWBORNS
BY BIRTHWEIGHT AND GESTATIONAL AGE



Preterm	Term	Postterm
---------	------	----------

ALBERTA



ALBERTA PRENATAL RECORD

This record is prepared and revised by the Perinatal Committee of the Alberta Medical Association, printed by the Department of Hospitals and Medical Care and distributed to the hospitals to be available to their doctors.

2 DOCTOR'S COPY
3 HOSPITAL COPY: MOTHER'S CHART
4 HOSPITAL COPY: NEWBORN CHART

A GUIDE TO PREGNANCY RISK

PREGNANCY AT HIGH RISK (10 times the average risk)

Pregnancies which are so complicated that the fetus and/or mother are obviously in danger. If at all possible, these patients should be transferred to a regional perinatal center for intensive care and delivery. Clearly, there are patients who deserve to be placed in this risk category who cannot be transferred safely or in time to benefit the fetus or mother.

Small for dates
Antepartum bleeding > 20 weeks
Twins (large for dates)
Polyhydramnios (Single fetus)
Insulin diabetes
Immaturity (31 weeks or less, > 20 weeks)

PREGNANCY AT RISK

The fetus and/or mother are definitely at risk and consultation should be obtained with a specialist obstetrician in your area. In addition, consultation with an appropriate internist may be necessary. These patients may be managed by continuing collaborative care and deliveries in an obstetrical unit with intermediate level nursing facilities OR they may be returned to the care of the referring physician with a suggested plan of management for the remainder of the pregnancy.

Gestational diabetes	> 42 weeks (postmaturity)
Uncertain dates	Preclampsia B.P. > 140/90, proteinuria
Rh. sensitization	Prolonged rupture of membranes
Smoking, alcohol, drugs	Incompetent cervix
Rubella	Chronic maternal conditions
Fever/amnionitis	(hypertension, heart, renal)
Abnormal presentation	

PREGNANCY AT LEAST RISK

Certain dates
No significant medical disease
No significant genetic history
No past pregnancy complications
No previous I.U.G.R. or perinatal mortality

SOME SUGGESTIONS AND INVESTIGATIONS

- Telephone consultation
- Sonar
- Symphysial fundal measurement
- Urinary estriols
- Fetal biophysical profile
- Fetal movement graphs
- Glucose Tolerance Test
- Non stress tests
- Fetal stress tests
- Rh. negative checks at 28, 32, 36 weeks
- X-ray
- Nitrazine speculum examination
- Cultures
- ?
- ?
- ?

PRENATAL RECORD

[illegible]

Exercise

DATE DISCONTINUED

	IN WEEKS		
UTERINE SIZE AT FIRST VISIT			EXPECTED
			ACTUAL

E.D.C.

[illegible]

Books

CURRENT MEDICATIONS

	NO	YES
EXCESS VOMITING	☐	☒
ALCOHOL	☐	☒
SMOKING	☐	☒
DRUG ABUSE	☐	☒

EXAMINATION: DATE

	NO	YES
DIABETES	☐	☒
TWINS	☐	☒
MALFORMATIONS	☐	☒
HEREDITARY DISEASE	☐	☒
OTHER	☐	☒

GENERAL
CONDITION

	NO	YES
RENAL	☐	☒
CARDIAC	☐	☒
INFECTIONS	☐	☒
INFERTILITY	☐	☒
ALLERGIES	☐	☒
OPERATIONS	☐	☒
TRANSFUSIONS	☐	☒
OTHER	☐	☒

HEAD & NECK

BREASTS

HEART

LUNGS

ABDOMEN

EXTREMITIES

BIMANUAL
VAGINAL

SPECULUM

PART II

PRENATAL RECORD

NAME

AGE	GRAV	PARA
-----	------	------

INITIAL INVESTIGATIONS

HEIGHT	PRE PREG WEIGHT	ABO	Rh	Hgb	SEROLOGY	RUBELLA TITRE	PAPP SMEAR	DATE OF QUICKENING	L N M P	E D C	
DATE	WGT.	URINE PRO / SUG	B.P.	GEST. AGE	FUNDUS (cm)	PRES. F.H.	REMARKS, MEDICATIONS, ETC.				NEXT VISIT

Topics for Discussion and Advice

Hosp. Adm. Procedures

Arrival at Hospital

Anaes. & Analg.

Labor Stages

Breast Care


Breast Feed:

Baby Care

RISK FACTORS

HIGH RISK INVESTIGATIONS

DATE					
------	--	--	--	--	--



NOTATIONS FOR DELIVERY/PUERPERIUM

PRENATAL CLASSES ☐

BREAST FEEDING ☐

BABY'S DOCTOR _____

DOCTOR'S SIGNATURE _____

NAME

[illegible]

SIGNED _____ M.D.

ALBERTA PRENATAL RECORD - Instructions

About fifty percent of perinatal deaths occur before the patient is admitted to hospital and it is likely that morbidity from premature birth and inadequate fetal growth can be reduced by more vigilant prenatal care.

This record has been prepared with the assistance of a widely representative group of physicians in Alberta in order to bring the recent advances in perinatal care into routine use throughout the Province and it is also hoped that the revised record will produce better carbon copies.

The two primary emphases at the first prenatal visit should be:

1. **ESTABLISHMENT OF E.D.C.**

To determine whether the patient is pregnant and, if so, by how many weeks. For this reason the data establishing the E.D.C. are listed first on the record and an attempt is made to arrange the pertinent information on the form in such a way that there is a logical sequence of thought and examination from the stated L.M.N.P. in the top left hand corner of this section to the final estimation of this important date in the bottom right hand corner of this section.

 - Menstrual Cycle** - Cycle length prior to L.M.P. (greater than 30 days) refers to menstrual pattern within three months of L.M.P.
 - Birth Control** - Oral contraceptives discontinued with one or two cycles before L.M.P.
- I.U.C.D. in situ may result in abortion or premature labor but may be removed during pregnancy. Prompt referral is desirable.
 - Bleeding** - Bleeding since L.M.P. refers to vaginal bleeding in early pregnancy. This can be associated with a molar pregnancy which, if suspected, would suggest estimation of urinary HCG and sonar examination.
 - Uncertain Dates** - Uncertain dates or lack of correlation between the period of gestation and the actual size of the uterus can lead to problems later in the pregnancy. Pelvic examination should include bi-manual palpation of the uterus.
 - Investigations to Confirm Dates** - Where dates are uncertain every effort should be made to establish gestational age accurately by sonar examination before the twentieth week of pregnancy (or uterine height of 20 cm.)
- The dates and results of pregnancy tests should be listed in this space.
- The Perinatal Committee has tried to avoid listing tests and investigations on this record in case this should be construed to mean that the tests were essential in all patients for good care.
2. **ASSESSMENT OF PATIENT'S HISTORIES AND GENERAL HEALTH TO FORECAST THE OUTCOME OF THIS PREGNANCY**

There is little change in this section of the prenatal record from the traditional approach to this assessment.

 - Obstetric History** - Only the most common recurrent complications are listed on the record but emphasis is placed on writing to the hospital for the case history if there is any doubt about previous adverse obstetric history or illness of the newborn.
 - Present Pregnancy** - The physiological changes in early pregnancy (nausea, breast changes, urinary frequency, etc.) have been omitted in this section but excessive vomiting requiring treatment should be noted and also smoking, alcohol and the use of non-medical drugs (eg. caffeine) as these lifestyles are known to affect the fetus adversely and are of importance in the care of the newborn. Every effort should be made to encourage the pregnant patient to change these three lifestyles because of the risk to the fetus.
 - Current Medications** - Any current prescription medication (steroids, insulin, tranquilizers) should be listed.
 - Family History** - This section has been shortened because a rather general question will usually illicit the pertinent information if the physician suspects there is history of which he should be aware. A history of seizures in the family or birth-damaged siblings, premature births, etc., would encourage a more detailed history.
 - Past Medical History** - This is not a complete list for the sake of brevity. For example, tuberculosis, venereal disease, and herpes, come collectively under "infections" but should be specified individually.

Physical Examination - The date of this examination should be specified even if it is the same as the date of first visit. Any abnormal findings or pelvic abnormalities should be listed.

Any risk factors identified on this first page should be noted in the RISK FACTOR section on the back of the form and, where necessary, the follow-up indicated e.g. "Dates uncertain" - Sonography (on such-and-such dates).

Repetition The back page of the record (Part II) has the patient's name, age, gravida and parity repeated at the top of the page. The usual ages considered at risk are less than 18 years and over 38 years but this depends on multiple factors. The primigravida over 30 years can cause concern. The para 4+ can prove a high risk patient.

Age Parity The initial investigations are listed at the top of the back page so that they are visible at every visit. If the reports on any of these investigations show deviations from the normal, e.g. Rh. antibodies, the follow-up would be in the high-risk boxes at the bottom of the page where the dates of follow-up tests would be planned and arrangements made regarding the father's genotype.

The initial investigations should be repeated at every pregnancy.

Hgb. The usual column for hemoglobin is omitted because the same recording procedure can be followed as for high risk deviations from the normal: a Hgb. of $<10g.$ would be checked off as a risk factor and listed in the high risk investigation boxes for treatment and further checks.

Height & Weight Height and weight are listed together but while women under 5 feet (152 cm.) and under 100 lbs. in weight (45 kgm.) are considered to be at risk, there is still a relevancy in family size (e.g. some immigrants are small people). It is the regular weight gain and the growth of the fetus that will be pertinent in the pregnancy and the weight column is directly below the pre-pregnancy weight for the purpose of watching weight gain. Obese women are at risk.

Date The first four columns are usually completed by the office nurse.

Urinalysis If the patient does not bring a specimen, the specimen should be obtained because this check is essential especially in the presence of even a slight increase in blood pressure. More than a trace of proteinuria warrants hospitalization.

Glycosuria in pregnancy should always suggest the presence of gestational diabetes until proved otherwise. Gestational diabetes requires prompt consultation.

Blood Pressure An increase in blood pressure (diastolic pressure increase of 20 mm. or a diastolic pressure of 90 mm.) requires careful assessment and possible hospitalization. Management of hypertension and/or edema in pregnancy with diuretic drugs on an ambulatory basis has now been proved ineffective because, while relieving the patient's symptoms, this treatment does not improve the basic placental dysfunction which can result in a small-for-dates or dead fetus.

Gestational Age Once the E.D.C. has been established the weeks of gestation of the pregnancy by dates should be entered on this column.

Fundus (cms.) Regardless of the gestational age by dates the height of the fundus should be carefully measured in centimetres at each prenatal visit, using a tape measure. The patient should lie flat with one pillow. The fundus is identified, without applying pressure to the abdomen, and the upper edge of the fundus is marked on the abdomen using a pen. The tape measure is then used to record the distance between the upper border of the symphysis pubis and the ink mark on the abdomen. This distance should be entered in the "Fundus" column.

Serial measurements of the size and rate of growth of the uterus are now considered so important in pregnancy that they warrant recording on the graph supplied in the bottom right hand corner of page two. This graph correlates weeks of gestation with the symphysis-fundal measurement. The expected recordings on this graph would be in zones 2 or 3. The dotted lines indicate the 10th and 90th percentiles and recordings above the 90th percentile at two successive visits would warrant a sonar scan (zone 1: twins, large infant, hydramnios, gestational age in doubt). On the other hand, if the uterus is small-for-dates (zone 4) or fails to increase in size between two visits there is a risk of feto-placental dysfunction and fetal growth retardation. This observation at two successive visits would warrant sonar examination and serial tests of feto-placental function after 28 weeks. Where there is inappropriate uterine growth for the gestational age (zone 1 or zone 4) there may also be less than normal fetal activity (less than ten movements during 12 hours on two successive days) and the patient should be asked to report this immediately.

Where intrauterine growth retardation is suspected serial sonar examination should be done (20, 26, 30, 34 and 38 weeks). The diameter of the fetal trunk should be estimated as well as the biparietal diameter as the ratio of this diameter to the fetal trunk diameter provides the most information in these cases.

The bottom right hand corner of the ultrasound chart requires the maternal abdominal length from the symphysis-pubis to the xiphoid. Women with short abdominal lengths (37 cm. or less) may have measurements recorded in the lower part of zone 3 or may even fall into zone 4 and give the appearance of being small-for-dates.

Presentation This column is for use after 34 weeks in case of a malpresentation. The column on "Engagement" was not considered essential and has been omitted.

F.H.R. This column is narrow and should only require a checkmark. Most doctors do not count the fetal heart beats in normal antenatal patients.

Topics for Discussion These are marginal reminder notes: the prenatal suggestions on the front page and the labor and delivery factors on the back page.

HIGH RISK FACTORS (Fly sheet in front) No attempt has been made to include complete lists because these lists are long and space-consuming. A longer check list could be supplied from the Perinatal office on request.

This half page with the risk factors on the back for constant reminder, may be used to conceal the name and maintain the confidentiality of records when they are in use during the prenatal period, or it may be detached if the doctor so wishes.



Saskatchewan
Health
Hospital Services
Plan

Prenatal Record

Health 19-42-5-79

Name		Age	Date of Birth
Address		Telephone No.	
Father of Child		Date of First Visit	
Family Physician	Obstetrician	Pediatrician	

Obstetrical History Including Abortions:								
Date	Place of Confinement	Dur. Preg. Wks.	Hours of Labour	Type of Delivery	Complications Mother and/or Infant	Child Sex	Birth Wt.	Pres. Health

Past illnesses: No Yes Specify	Family history: No Yes Specify
Renal	Diabetes
Cardiac	Cardiac
Infections	Hypertension
Rubella	Tuberculosis
Venereal	Twins
Allergies	Malformations
Operations	Mental Retardation
Transfusions	Other
Other	

Menstrual history: Cycle _____ Duration _____ L M P _____ Normal Abnormal	History of present pregnancy: Specify	Current medications
Method of contraception: Date of Discontinuance _____	Nausea, Vomiting No Yes Bleeding No Yes Date Fever No Yes Date Radiation No Yes Date Smoking No Yes Amount Drug Use No Yes Identify Nutrition Poor Good	
Pre-Preg. Weight _____ Height _____		

Examination Date _____	
General Condition: _____	Breasts: _____
Head & Neck: _____	Abdomen: _____
Lymph Nodes: _____	Spine: _____
Heart: _____	Limbs: _____
Lungs: _____	Pelvic Examination: _____

Fetal Risk Scoring "A" Baseline Data (Maximum Score is 3)		Previous Obstetrical History
Age 35 + 1 ☐	Rh iso - Immunized Mother 2 ☐	Abortion ☐
40 + 2 ☐	Pos. History of Erythroblastosis 3 ☐	Stillbirth ☐
Para 0 1 ☐		Neonatal Death ☐
6 + 2 ☐		Surviving Premature Infant ☐
Interval less than 2 years 1 ☐		Antepartum Hemorrhage ☐
Weight 200 lb. + 1 ☐		Toxemia ☐
Diabetes - Mild, Moderate 2 ☐		Mid-Forceps Delivery ☐
Severe 3 ☐		Cesarian Section ☐
Chronic Renal Disease 1 ☐		Major Congenital Anomaly ☐
with Diminished Renal Function 3 ☐		Baby 10 lbs. + ☐
Pre-Existing Hypertension 1 ☐		One Instance of Above 1 ☐
140 + /90 + 1 ☐		Two or More Instances of Above 2 ☐
160 + /110 + 2 ☐		

Total Score Part "A"

Doctor's Chart

[illegible]

PROVINCE OF MANITOBA

OBSTETRICAL RECORD

NAME		ADDRESS		PHONE		DATE	
DOCTOR		HOSPITAL		INSURANCE-M.B.S.C.		WARD	
AGE		MENSTRUATION		PILL		L.H.B.P.	
E.S.C.		YEAR OF BIRTH		HOSPITAL NUMBER		PHYSICIAN	
☐ 37 YEARS OR GREATER AMNIOCENTESIS ☐ DONE ☐ NOT DONE ☐ REFUSED							
OBSTETRICAL <u>GRAV.</u> <u>PARA</u> <u>ALIVE</u>							
PREMATURE OR < 3200 gm ☐ ALIVE ☐ DEAD STILLBIRTHS							
YEAR		WT.		GEST.		COMPLICATIONS	
POST PARTUM		HIGH RISK SCORE		1ST VISIT		2ND VISIT	
PREV. C/S		ALLERGIES - DRUGS		ETHNIC GROUP		PREV. TRANSFUSION	
FAMILY HX		CONGENITAL DEAFNESS		TB		DIABETES	
GENETIC HX		TWINS		HYPERTENSION		BLEEDING	
MED. AND SURG.		DIABETES		CORE BS		EPILEPSY	
PRESENT HX		MEDICATION		INFECTION		INFERTILITY	
EXAMINATION		DATE		WEIGHT		B.P.	
HEAD AND NECK		BREASTS		PELVIC		HEIGHT	
HEART		LUNGS		CERVIX		S.T.S.	
ABDOMEN		EXTREMITIES		CORPUS		Rh.	
OBSTETRICAL PELVIC ASSESSMENT		ADNEXA		LAB. NO.		IMMUNOGLOBULIN	
URINE		P		S		MICRO	
CERVIX		PAP.		C AND S		BREVINDEX	
DATE		WT.		BP		URINE	
WEEKS		FUNDUS		POSITION		FM	
HOB AND REMARKS		RETURN		ID		RUBELLA H.I.T.	
BLOOD SUGAR		YES		NO		ANTIBODIES	

PRENATAL SCORING FORM

1) Score each question as indicated 2) Total each category score at 1st visit 3) Repeat at 36 weeks 4) Repeat on admission 5) Record on P N sheet

I. REPRODUCTIVE HISTORY		II. ASSOCIATED CONDITIONS		III. PRESENT PREGNANCY			
				1st VISIT	36 WEEKS	ON ADM	
AGE	$<16 = 1$ $16 - 35 = 0$ $>35 = 2$	PREVIOUS GYNAECOLOGICAL SURGERY	= 1 ☐	BLEEDING			
PARITY	$0 = 1$ $1 - 4 = 0$ $5 + = 2$	CHRONIC RENAL DISEASE	= 2 ☐	<20 WEEKS	= 1 ☐	☐	☐
PAST OB HISTORY		GESTATIONAL DIABETES	= 1 ☐	>20 WEEKS	= 3 ☐	☐	☐
HABITUAL ABORTION/ INFERTILITY	= 1 ☐	DIABETES MELLITUS	= 3 ☐	ANEMIA <10 gm %	= 1 ☐	☐	☐
P.P.H/MANUAL REMOVAL	= 1 ☐	CARDIAC DISEASE	= 3 ☐	PROLONGED PREGNANCY (42 WEEKS)	= 1 ☐	☐	☐
BABY >9 LBS (4082 gm)	= 1 ☐	OTHER MEDICAL DISORDERS		HYPERTENSION	= 2 ☐	☐	☐
BABY <5½ LBS (2500 gm)	= 1 ☐	(Chronic bronchitis, Lupus, Etc)		PREMATURE RUPTURE MEMBRANES	= 2 ☐	☐	☐
P.E.T./HYPERTENSION	= 1 ☐	SCORE ACCORDING TO SEVERITY = ☐		POLYHYDRAMNIOS	= 2 ☐	☐	☐
PREVIOUS CAESAREAN	= 2 ☐	(1 to 3)		SMALL FOR DATES	= 3 ☐	☐	☐
STILLBIRTH OR NEONATAL DEATH	= 3 ☐			MULTIPLE PREGNANCY BREECH	= 3 ☐	☐	☐
PROLONGED LABOUR OR DIFFICULT DELIVERY	= 1 ☐			MALPRESENTATION			
				Rh ISOIMMUNIZATION	= 3 ☐	☐	☐
CATEGORY SCORE _____ +		CATEGORY SCORE _____ +		CATEGORY SCORE _____			

TOTAL RISK SCORE

1st VISIT = ☐

TOTAL AT 36 WEEKS = ☐

TOTAL ON ADMISSION TO HOSPITAL = ☐

(Record scores in box on prenatal sheet)

NOTE: LOW RISK = 0 - 2

HIGH RISK = 3 - 6

EXTREME RISK = 7 >

COMMENTS

NOTES

DELIVERY RECORD	TYPE OF LABOUR	INDUCTION	ANAESTHESIA
DATE OF BIRTH	SPONTANEOUS	FORCEPS	THIRD STAGE
HOUR OF BIRTH	WEIGHT	L.B./S.B./N.N.D.	EPISIOTOMY
PRES. POS.	BLOOD LOSS		NURSING
BABY - SEX			
COMPLICATIONS			

POST PARTUM EXAMINATION	DATE	DOCTOR'S SIGNATURE	
WEIGHT	LOCHIA		
S.P.	MENSES		
HAEMOGLOBIN	NURSING	PELVIS	PERINEUM
URINE	BREASTS	CERVIX	CORPUS
FAMILY PLANNING			

HOSPITAL _____

**CANARY - MOTHER'S CHART
FORWARD TO HOSPITAL**

[illegible]


PRESS HARD
THIS FORM IS IN
TRIPLICATE

WHITE - INFANT'S CHART

PINK - DOCTOR'S COPY

CANARY - MOTHER'S CHART

PATIENT'S NAME	
ADDRESS	
DOCTORS	
BANK FACTORS IDENTIFIED BY FACTA 1957	
	QUICKENING

[illegible]**ADDITIONAL COMMENTS:**

ULTRASOUND AND ABNORMAL LABORATORY RESULTS

	NUTRITION
	DIET
	REST
	TRAVEL
	PREM. LABOUR
	BREAST CARE
	ANAES MED.
	TYPE OF DELIVERY
	FATHER AT DELIVERY
	BREAST FEEDING
	ROOM-IN
	CIRCUMCISION
	EARLY DISCHARGE
	CAR SAFETY
	OBSTETRICIAN
	PAEDIATRICIAN

MEASURE SYMPHYSIS-FUNDUS
HEIGHT AT EACH ANTENATAL
VISIT AND RECORD ON GRAPH
OPPOSITE.

PHYSICIAN'S SIGNATURE _____

POSTNATAL VISIT

DATE _____

WEEKS POSTPARTUM: _____

BREAST FEEDING

☐ YES

☐ NO

REASONS FOR DISCONTINUING _____

PHYSICAL EXAM

WT _____ kg.

B P _____ mmHg.

EPISIOTOMY : _____ C. SECTION INCISION: _____

RUBELLA IMMUNE

☐ YES

☐ NO

VACCINATED

☐

URINALYSIS: _____

PAP: _____

UNRESOLVED PROBLEMS: _____

ADVICE RE FUTURE PREGNANCIES (EG. CONTRACEPTION, RISK STATUS, REQUIREMENT FOR ULTRASOUND ETC.)

PHYSICIAN'S SIGNATURE _____

A GUIDE TO PREGNANCY RISK GRADING

At each antenatal visit please give your assessment of pregnancy (fetal plus maternal) risk according to the following grading system. The risk factors or problems listed below are intended as *examples only*. Additional space is provided for other risk producing problems which you have identified. This risk grading system is intended as a basis for planning the ongoing management of the pregnancy.

Pregnancy at no predictable risk

- ☐ NO PRIOR PERINATAL MORTALITY OR LOW BIRTHWEIGHT INFANT
- ☐ NO SIGNIFICANT MEDICAL DISEASE
- ☐ NO PREGNANCY COMPLICATIONS NOW OR IN THE PAST (Bleeding, Hypertension, Premature Labour)
- ☐ FETAL GROWTH SEEMS ADEQUATE

Pregnancy at risk

The fetus and/or mother may be at risk and consultation should be considered with a specialist obstetrician in your area. In addition, consultation with an appropriate internist may also be necessary. (These patients may be managed by continuing collaborative care and delivery in an obstetrical unit with intermediate level nursing facilities OR they may be returned to the care of the referring physician with a suggested plan of management for the remainder of the pregnancy).

- | | |
|---|---|
| ☐ DIABETES, CLASS A (GESTATIONAL) OR CLASS B | ☐ RENAL DISEASE WITHOUT HYPERTENSION |
| ☐ HYPERTENSION WITHOUT TOXAEMIA | ☐ MILD TOXAEMIA |
| ☐ APH, CEASED AND IN HOSPITAL | ☐ CONTROLLED PREMATURE LABOUR |
| ☐ CERVICAL INCOMPETENCE | ☐ MULTIPLE PREGNANCY |
| ☐ HYDRAMNIOS | ☐ BREECH: BEYOND 36 WEEKS |
| ☐ POST - DATE PREGNANCY (42 weeks +) | ☐ PRIMIGRAVIDA (age 35 +) |
| ☐ HISTORY OF PRIOR STILL BIRTH OR NEONATAL DEATH OR PREMATURE DELIVERY | ☐ HISTORY OF GENETIC DISEASE IN FAMILY (Genetic Amniocentesis or Counselling required) |
| ☐ MATERNAL OBESITY (20% > IDEAL B.WT.) | ☐ ANAEMIC NOT RESPONDING TO IRON (< 10gm) |
| ☐ SIGNIFICANT TOBACCO, ALCOHOL, DRUG INTAKE | ☐ WEIGHT GAIN < 10 lbs. BY 30 WEEKS |
| ☐ RHESUS IMMUNIZATION | ☐ GRAND MULTIPARA (> 5) |
| ☐ PREMATURE RUPTURE OF MEMBRANES 34 WK. OR MORE | ☐ LARGE FOR DATES |
| ☐ OTHER SIGNIFICANT MEDICAL ILLNESS | ☐ PREVIOUS UTERINE SURGERY INCLUDING CAESAREAN SECTION |
| | ☐ |

Pregnancy at high risk

Pregnancies which are so complicated that the fetus and/or mother are obviously in danger. If at all possible, these patients should be transferred to a regional perinatal centre (level III) for intensive care and delivery. Clearly, there are patients who deserve to be placed in this risk category (with problems such as excessive antepartum bleeding, cord prolapse, or advanced uncontrolled premature labour) who cannot be transferred safely or in time to benefit the fetus or mother.

- | | |
|---|--|
| ☐ DIABETES CLASS C, D, F, R OR SIGNIFICANTLY COMPLICATED | ☐ HYPERTENSION WITH SUPERIMPOSED TOXAEMIA |
| ☐ RENAL DISEASE WITH HYPERTENSION ± ↓FUNCTION | ☐ EARLY UNCONTROLLED PREMATURE LABOUR * |
| ☐ PREMATURE RUPTURE OF MEMBRANES (± SEPSIS) * | ☐ SEVERE FETAL GROWTH ARREST (< 10th Percentile) |
| ☐ ANTEPARTUM BLEEDING, CONTINUING OR REPEATED * | ☐ HEART DISEASE, ESPECIALLY WITH FAILURE |
| ☐ | ☐ |
| ☐ | ☐ |

* 24 - 34 weeks gestational age.

Two or more minor risk problems can combine to produce a high pregnancy risk. Such a patient may deserve to be placed in a higher risk category.

DOSSIER PRÉNATAL No 1

HÔPITAL		SOINS OBS.		SOINS BÉBÉ	
MÉDECIN					
NOM DE LA MALADE			RELIGION		
ADRESSE					
DATE DE NAISSANCE		J / M / A	ÂGE		TÉL.
PROFESSION		ORIGINE ETHNIQUE		ÉTAT CIVIL	
NIVEAU D'INSTRUCTION		OHIP NO.			
NOM DU PÈRE		DATE DE NAISSANCE		ÂGE	PROF.
SITUATION FAMILIALE					

Cycle D.M.: Normale D.M (date)		Durée Anormale	Orale Durée (années) Stérilet	Amét (date)	Dates J / M / A Dimensions	D.P.A. revisée J / M / A
--------------------------------------	--	-------------------	-------------------------------------	-------------	----------------------------------	-----------------------------

ANAMNÈSE MÉDICALE	NUTRITION	ANTÉCÉDENTS GÉNÉRAUX	ANTÉCÉDENTS GYNÉCOLOGIQUES

MAL REINS	HYPERTENSION	PERTES DE SANG
HYPERTENSION	MAL COEUR	PYREXIE
INFECTIONS	MALFORMATION	ALCOOL
MAL THYROÏDE	MAL GÉNÉTIQUES	MÉDICAMENTS
OPÉRATIONS		ALLERGIES

PRODUITS LAITIERS	GAIN POIDS
FRUITS ET LÉGUMES	ALCOOL
ALLAITEMENT	PLANS TRAY
COIT	

COCHER (-) SI NORMAL	
POITRINE	O.R.L.
TÊTE, COU, THYROÏDE	VULVE
ABDOMEN	COL
VARICES	OVAIRES

DATE DE LA PREMIÈRE VISITE	J / M / A	SIGNATURE DU MÉDECIN
----------------------------	-----------	----------------------

DOSSIER PRÉNATAL N° 2

MÉDECIN

NOM DU MALADE

ADRESSE

DATE DE DÉBUT DE LA GROSSESSE

DATE

J / M / A

M / A

PROGÉNÉDURE

PROGÉNÉDURE

PROGÉNÉDURE

RÉSULTAT D'ÉVALUATION


PROGÉNÉDURE

APRÈS FERMETURE
Poursuite en 3 parties

U.B.E.R. - NOÛ

FEUILLE DE LA MÈRE - BLANCHE: FEUILLE DU MÉDECIN - JAUNE: FEU

À CHAQUE VISITE PRÉNATALE, INSCRIRE LES CATÉGORIES COURANTES SUSMENTIONNÉES DE RISQUES DE GROSSESSE — A, B ou C — UTILISER CETTE CATÉGORIE POUR LE TRAITEMENT CONTINU DE LA GROSSESSE.
par ex.: C ENVISAGER LE TRANSFERT
B ENVISAGER UNE CONSULTATION
DEUX RISQUES MINEURS OU PLUS COMBINÉS PEUVENT PRODUIRE UN RISQUE SÉRIEUR DE GROSSESSE



MESURER LA HAUTEUR DE SYMPHYSE-FOND DE L'UTÉRUS À CHAQUE VISITE PRÉNATALE ET L'INSCRIRE SUR LE GRAPHIQUE CI-CONTRE.

NUTRITION	
REPOS	
COURS PRÉNATAUX	
TYPE D'ACCOUCHEMENT	
ALLAITEMENT	
CONGÉ PRÉCOCE	
CONTRÔLE DES NAISSANCES	
PÉDIATRE	

SIGNATURE DU MÉDECIN

GUIDE DE GRADATION DES RISQUES DE GROSSESSE

A chaque visite prénatale, indiquez votre évaluation des risques de la grossesse (foetus et mère) selon l'échelle de gradation suivante. Les facteurs de risques ou les problèmes énumérés ci-dessous ne sont que des *exemples*. Il reste des blancs où vous pouvez inscrire d'autres problèmes comportant des risques que vous avez relevés. Cette échelle de gradation des risques devrait servir de base à la planification du traitement continu de la grossesse.

CATÉGORIE A Grossesse sans risque prévisible

- ☐ AUCUNE MORTALITÉ PÉRINATALE ANTÉRIEURE OU BÉBÉ DE POIDS INFÉRIEUR
- ☐ AUCUNE MALADIE MÉDICALE IMPORTANTE
- ☐ AUCUNE COMPLICATION DE GROSSESSE MAINTENANT OU AUPARAVANT (Perte de sang, hypertension, travail prématuré)
- ☐ CROISSANCE APPAREMMENT NORMALE DU FOETUS

CATÉGORIE B Grossesse comportant des risques

Le foetus et (ou) la mère sont définitivement susceptibles à des complications et il faut obtenir une consultation d'un spécialiste en obstétrique de votre région. De plus, il faudra peut-être une consultation auprès d'un interniste de la spécialité voulue. On peut traiter ces cas grâce à des soins coopératifs continus dans une unité d'obstétrique où se fera l'accouchement et où existent des installations de soins infirmiers de niveau intermédiaire OU on peut les remettre entre les mains du médecin traitant en lui proposant une conduite à suivre applicable au reste de la grossesse.

- | | |
|---|--|
| ☐ DIABÈTE (CATÉGORIE A (GESTATIONNEL) OU CATÉGORIE B | ☐ AFFECTION RÉNALE SANS HYPERTENSION |
| ☐ HYPERTENSION SANS TOXÉMIE | ☐ TOXÉMIE BÉNIGNE |
| ☐ HAP, TERMINE ET À L'HÔPITAL | ☐ TRAVAIL PRÉMATURÉ CONTRÔLÉ |
| ☐ INCOMPÉTENCE CERVICALE | ☐ GROSSESSE MULTIPLE |
| ☐ HYDRAMNIOS | ☐ PRÉSENTATION PAR LE SIEGE |
| ☐ GROSSESSE PASSE TERME (42 semaines +) | ☐ PRIMIGESTE (âge 35 +) |
| ☐ ANTECÉDENTS DE MORTINAISSANCE OU DE DÉCÈS NÉONATAL | ☐ ANTECÉDENTS FAMILIAUX DE TROUBLES GÉNÉTIQUES (Amniocentèse génétique ou besoin de consultation) |
| ☐ OBESITÉ MATERNELLE | ☐ ANÉMIE FERRIPRIVE RÉFRACTAIRE |
| ☐ USAGE IMPORTANT DE TABAC, D'ALCOOL, DE DROGUES | ☐ GAIN DE POIDS < 10 liv. à 30 semaines |
| ☐ IMMUNISATION AU RHÉSUS | ☐ GRANDE MULTIPARE |
| ☐ | ☐ |
| ☐ | ☐ |

CATÉGORIE C Grossesse à risque élevé

Grossesses si compliquées que le foetus et (ou) la mère sont manifestement en danger. Si cela est possible, on devrait transférer ces cas à un centre périnatal régional (niveau III) pour soins intensifs et accouchement. De toute évidence, il y a des cas qui méritent d'être classés dans cette catégorie de risques (problèmes de pertes de sang excessives avant parturition, prolapsus du cordon ou travail prématuré avancé non contrôlé) mais qu'on ne peut transférer en sécurité ou à temps pour que ce soit avantageux au foetus ou à la mère.

- | | |
|---|---|
| ☐ DIABÈTE, CATÉGORIE C, D, F, R OU SUFFISAMMENT COMPLIQUÉ | ☐ HYPERTENSION COMPLIQUÉE DE TOXÉMIE |
| ☐ AFFECTION RÉNALE ACCOMPAGNÉE D'HYPERTENSION $\pm$ FONCTION | ☐ TRAVAIL PRÉMATURÉ NON CONTRÔLÉ* |
| ☐ RUPTURE PRÉMATURÉE DE LA POCHÉ DES EAUX ($\pm$ SEPTICÉMIE)* | ☐ ARRÊT DE CROISSANCE IMPORTANT DU FOETUS (< 10 ^e percentile) |
| ☐ PERTES DE SANG AVANT PARTURITION, CONTINUES OU REPÉTÉES* | ☐ TROUBLES CARDIAQUES, SURTOUT AVEC DÉFAILLANCE |
| ☐ | ☐ |
| ☐ | ☐ |

* en particulier 24 à 34 semaines de gestation

La présence de deux risques mineurs ou plus peut produire une grossesse à risque élevé.
Il se peut que ces cas méritent d'être dans une catégorie à risque plus élevé.

DOSSIER OBSTÉTRICAL
OBSERVATION MÉDICALE

Cadre réservé à l'usage du centre hospitalier

Nom _____

Prénom _____

Adresse _____

_____ N° de téléphone _____

Âge _____ Date de naissance _____

N° d'assurance-maladie _____

Service de consultation	Service de chirurgie	Service de gynécologie
Service de pédiatrie	Service de médecine interne	Service de dermatologie
Service de radiologie	Service de laboratoire	Service de pharmacie
Service de nutrition	Service de psychiatrie	Service de réhabilitation
Service de soins infirmiers	Service de soins dentaires	Service de soins de suite

ANTÉCÉDENTS FAMILIAUX

Tuberculose, hypertension, cardiopathie, diabète, allergie, cancer, troubles neuro-psychiatriques, grossesse gémellaire, anomalies.

Préciser:

ANTÉCÉDENTS MENSTRUELS

Première menstruation	Cycle	Durée	D.D.M.	D.P.A.	Méthode anticonceptionnelle	Cessée depuis
ans	jours	jours				

ANTÉCÉDENTS PERSONNELS (✓ : positif O : négatif)

Antécédents éloignés			Antécédents récents (depuis les dernières menstruations)		
Maladies rénales		Phlébites, varices	Nausées		Rougeole — Varicelle
Maladies cardiaques		Épilepsie	Vomissements		Autres infections virales
Hypertension artérielle		Allergies aux médicaments	Mauvaise digestion		Irradiations, radiographie
Rhumatisme articulaire aigu		Allergie	Constipation		Traumatismes
Tuberculose		Dyscrasies sanguines	Céphalées		Médicaments
Maladies vénériennes		Incompatibilité Rh ABO	Hémorragies (préciser)		Vaccins
Maladies gynécologiques		Opérations antérieures	Pertes vaginales (préciser)		Drogues
Infections virales		Accidents antérieurs	Oedème		Alcool
Maladies neuro-psychiatriques		Vaccins	Transfusions de sang		Cigarette
Diabète		Génétique	Douleurs abdominales (préciser)		Travail à l'extérieur
Affections de la thyroïde		Immunoglobuline anti-D	Symptômes urinaires		

Préciser:

Grossesses antérieures			À terme	Prématurées	Dysmatures	Avortements	Enfants vivants	Grossesse multiple	Grossesse ectopique
N°	Date	Centre hospitalier	Durée de la grossesse	Durée du travail	Incidents de l'accouchement	♂ ou ♀	Masse (poids)	Chez la mère	Chez l'enfant

DOSSIER OBSTÉTRICAL
OBSERVATION MÉDICALE

Cadre réservé à l'usage du centre hospitalier

Nom _____
Prénom _____
Adresse _____
N° de téléphone _____
Âge _____ Date de naissance _____
N° d'assurance-maladie _____

Prénom	N° de naissance	N° du dossier médical
Prénom	N° de naissance	N° du dossier médical
Prénom	N° de naissance	N° du dossier médical
Prénom	N° de naissance	N° du dossier médical
Prénom	N° de naissance	N° du dossier médical
Prénom	N° de naissance	N° du dossier médical
Prénom	N° de naissance	N° du dossier médical
Prénom	N° de naissance	N° du dossier médical
Prénom	N° de naissance	N° du dossier médical
Prénom	N° de naissance	N° du dossier médical

EXAMEN PHYSIQUE			
Taille	m	Masse (poids) avant la conception	kg
Masse (poids) actuelle	kg		
Yeux	Dents	Pharynx	Thyroïde
Cou	Peau	Cœur	Poumons
Seins	Mamelons	Tumeurs	
Abdomen		Colonne	
Membres inférieurs	Oedème	Varices	
EXAMEN GYNÉCOLOGIQUE			
Vulve			
Vagin			
Périnée			
Col			
Utérus			
Correspondance entre le volume de l'utérus et l'âge de la grossesse			
Annexes			
Toucher rectal		Cul-de-sac de Douglas	
Pelvimétrie	Clinique	Promontoire	Épines sciatiques
	Radiologique	Oui ☐ Non ☐	Si oui, date
Dystocie prévue au détroit		supérieur ☐	moyen ☐ inférieur ☐
Cytologie	Oui ☐ Non ☐	Si oui, date	
Problèmes spéciaux			
Type d'accouchement prévu		Type d'anesthésie prévu	Allaitement maternel prévu

_____ Date

MEDECIN TRAITANT

_____ Signature

2

Nom à la naissance

Prénom(s) _____

Adresse _____

N° de téléphone _____

Âge _____ Date de naissance _____

N° d'assurance-maladie _____

Norm du conjoint _____

EXAMENS DE LABORATOIRE

B.W. _____ Date _____

Hb _____ Ht _____ Date _____

Groupe et Rh de la mère	Anticorps	Cytologie
-------------------------	-----------	-----------

Urines (sommaire et culture) _____

D.P.A. selon:

1.0 D.M.

2. clinique

3. échographie

2

DOSSIER OBSTÉTRICAL
ÉVOLUTION DU TRAVAIL

Nom de l'établissement _____

Date d'admission _____ heure _____ h _____

Date du début du travail _____ heure _____ h _____

Membranes ☐ intactes ☐ rompues ☐ h _____ liquide _____

État de la parturiente à son admission

☐ Passage	Par.
☐ Levain	Ab
☐ Glycolurie	Gr.
☐ Albuminurie	D.M.
☐ Hémogramme et groupe sanguin	Sem. de gestation
☐ Culture d'urine	D.P.A.

Médecin prévenu à _____ h _____

[illegible]


DOSSIER OBSTÉTRICAL

COURBE DE FRIEDMAN (1)

Nom de l'établissement _____

Raison d'admission

- ☐ Membranes rompues ☐ Travail vrai ☐ Césarienne non urgente
☐ Suspension du travail ☐ Autre _____
☐ Indication du déclenchement _____



(1) Friedman, Emmanuel A. Labor in Multiparas. Extrait de: Obst. & Gynec. vol. 8 n° 6; 1956. p. 696.

Date	Heure	Médication (inscrire la date, l'heure, le nom, la quantité et la voie d'administration)	Initiales
	h		
	h		
	h		
	h		
	h		
	h		
	h		
	h		
	h		
	h		
	h		
	h		
	h		

DOSSIER OBSTÉTRICAL
GROSSESSE, TRAVAIL ET ACCOUCHEMENT
ÉVOLUTION DE LA MÈRE ET
ÉVALUATION DU NOUVEAU-NÉ

Cadre réservé à l'usage du centre hospitalier

Nom de l'établissement _____

GROSSESSE, TRAVAIL ET ACCOUCHEMENT				
D.D.M.		a	m	j
D.P.A.		a	m	j
Groupe et Rh				
Par:	Gr.	Ab.	Enfants vivants	Particularités (complications ou diagnostics pendant cette grossesse ou les grossesses antérieures)
Anticorps				
Présentation				
☐ tête ☐ siège ☐ autre (préciser) _____				
Travail				
☐ Spontané ☐ Déclenché				
Début du travail		a	m	j
1er stade		durée _____		
2e stade		durée _____		
3e stade		durée _____		
Membranes rompues		a	m	j
durée totale		_____ h		
Analgésie durant les 4 dernières heures (médicament, dosage)				
Cortico-stéroïdes (date et heure de la première dose)				
Anesthésie				
☐ Aucune ☐ Générale ☐ Bloc para-cervical ☐ Epidurale ☐ Bloc honteux ☐ Locale				
Accouchement				
a		m	j	h
☐ Par voie vaginale ☐ Par césarienne				
☐ TÊTE ☐ SIÈGE				
☐ Spontané ☐ Spontané ☐ Itérative ☐ Forceps bas ☐ Assisté ☐ Transversale basse ☐ Forceps moyens ☐ Forceps ☐ Médiane basse ☐ Rotation ☐ Classique ☐ Hystérectomie				
Indication de la césarienne				

Épisiotomie		Aucune	Médiane	Médo-latérale
		☐	☐	☐
Déchirure		Aucune	Périnéale 1 2 3 4	Cervicale
		☐	☐	☐
		Vaginale		
		Pertes sanguines _____ ml		
Cordon ombilical		Circulaires		
Clampage		☐ Lâches		
☐ < 45 secondes après l'accouchement		☐ Serrés		
☐ > 45 secondes après l'accouchement		☐ Coupés pendant l'accouchement		
		☐ Coupés après l'accouchement		
Placenta		Anomalies		
Masse _____ g		☐ Évacuation spontanée		
		☐ Évacuation manuelle		
		☐ Vaisseaux ombilicaux 2 3		
Complications		Aucune	Meconium	Détresse fœtale
		☐	☐	☐
☐ < 120		☐ clinique		
Coeur fœtal ☐ 120 à 160		☐ Monitoring		
☐ > 160		☐ externe		
		☐ interne		
Signature du médecin		Date		

ÉVALUATION DU NOUVEAU-NÉ				
Sexe	☐ ♂ ☐ ♀ ☐ Vivant ☐ Mort-né	Masse _____ g		
☐ Gouttes ophtalmiques		☐ Vitamine K	☐ Ondoyé	
Alimentation		☐ Maternelle	☐ Artificielle	☐ homo. ☐ humanisé
Réanimation	Ventilation			
☐ oui ☐ non	☐ Avec sac et masque ☐ Tube trachéal _____ min.	☐ Aspiration trachéale		
Asphyxie		☐ Aucune	☐ Modérée	☐ Importante
☐ Anomalies		Spécifier: _____		
☐ Complications		Parents informés ☐ oui ☐ non		
APGAR		0	1	2
Battements cardiaques		Absents	Moins de 100	Plus de 100
Efforts respiratoires		Absents	Lents irréguliers	Bons pleurs
Tonus musculaire		Fiasque	Flexion des extrémités	Mouvements actifs
Réflexes à la stimulation		Absents	Grimace	Pleure avec force
Coloration des téguments		Bleue pâle	Corps rose extrémités bleues	Entièrement rose
Remarques		Total		
Nom du médecin qui suit l'enfant		Date		

ÉVOLUTION DE LA MÈRE				
Suites de couche		Hémorragie des suites de couche	Immunoglobuline anti-D	Fièvre ☐ oui ☐ non
☐ Normales		☐ Immédiate	Donnée le	☐ Endométrite
☐ Anormales		☐ Tardive	a	☐ Autre infection pelvienne
Hb la plus basse _____		Transfusion _____ sacs	m	☐ Trombo-embolie
Dernière Hb _____			j	☐ Infection urinaire
État à la sortie		Vivante	Décédée	☐ Infection respiratoire
Date		☐	☐	Signature du médecin
		☐	☐	Date

OBSTETRICAL CHART
MEDICAL HISTORY

Space reserved for the use of the hospital center

Surname _____
 Given name(s) _____
 Address _____
 _____ Telephone No. _____
 Age _____ Date of birth _____
 Health insurance No. _____
 Name of spouse _____

PHYSICAL EXAMINATION									
Height (meters) _____ m			Weight before pregnancy _____ Kg			Weight now _____ Kg			
Eyes		Teeth		Pharynx			Thyroid		
Neck		Skin		Heart			Lungs		
Breasts			Nipples			Tumours			
Abdomen						Spine			
Lower limbs			Edema			Varicose veins			
GYNECOLOGICAL EXAMINATION									
Vulva									
Vagina									
Perineum									
Cervix									
Uterus									
						Relationship between size of uterus and weeks of gestation			
Annexae									
Rectal									
						Pouch of Douglas			
Pelvimetry		Clinical	Promontary			Ischial spines			
		Radiological	Yes ☐	No ☐	If yes, date	yr	m	day	
Dystocia anticipated		Inlet		Midplane		Outlet			
		☐		☐		☐			
Cytology		Yes ☐	No ☐	If yes, date		yr	m	day	
Special problems									
Type of delivery anticipated				Type of anaesthesia anticipated			Breast feeding anticipated		

Space reserved for the use of the hospital center

Name of spouse _____

Urine (summary and culture) _____

E.D.C. according to	y:	m	day
1. L.M.P.			
2. Clinical			
3. Echography			

[illegible]

Name of establishment _____

Date of admission _____ Hour _____ h _____

Date of onset of labour _____ Hour _____ h _____

☐ Spontaneous ☐ Induced

Membranes ☐ Intact ☐ Ruptured _____ h _____ Amniotic fluid _____

STATE OF PATIENT UPON ADMISSION

STATE OF PATIENT UPON ADMISSION

☐ Shave prep.	Para
☐ Enema	AB
☐ Glycosuria	GR
☐ Albuminuria	D.M.:
☐ Hemogram and blood group	Weeks of gestation
☐ Urine culture	E.D.C.

Physician notified at _____ h _____

Physician notified at _____ h _____


[illegible]

OBSTETRICAL CHART FRIEDMAN'S CURVE

Name of establishment _____

Reason for admission

- ☐ Ruptured membranes ☐ True labour ☐ Elective c. section
- ☐ False labour ☐ Other _____
- ☐ Induction _____



Date	Time	Medication (indicate date, time, name, dosage, and route of administration)	Initials
	h		
	h		
	h		
	h		
	h		
	h		
	h		
	h		
	h		
	h		
	h		
	h		
	h		

5

OBSTETRICAL CHART
PREGNANCY, LABOUR AND DELIVERY
EVOLUTION OF THE MOTHER AND
EVALUATION OF THE NEWBORN

Space reserved for the use of the hospital center

Name of establishment _____

PREGNANCY, LABOUR AND DELIVERY									
L.M.P. yr m day			E.D.C. yr m day			Group and Rh			
Par	Gr	Ab	Live children	Peculiarities (complications or diagnostics since this pregnancy or past pregnancies)					
Presentation ☐ Vertex ☐ Breech ☐ Other (specify) _____									
Labour ☐ Spontaneous ☐ Induced Onset of labour yr m day _____ h _____ 1st stage duration _____ 2nd stage duration _____ 3rd stage duration _____ Total duration _____ Membranes ruptured yr m day _____ h _____									
Anaesthesia during last 4 hours (medicines, dosage)									
Corticosteroids (date and hour of 1st dosage)									
Anaesthesia None ☐ General ☐ Para-cervical ☐ Epidural ☐ Pudental ☐ Local ☐									
Delivery yr m day _____ h _____ ☐ Vaginal ☐ Caesarian section ☐ VERTEX ☐ BREECH ☐ Spontaneous ☐ Spontaneous ☐ Low forceps ☐ Assisted ☐ Mild forceps ☐ Forceps ☐ Rotation ☐ Iterative ☐ Low transverse indication ☐ Low median ☐ Classical ☐ Hysterectomy									
Indication of caesarian section									

Episiotomy None ☐ Median ☐ Medio-lateral ☐			
Laceration None ☐ Perineal 1 2 3 4 ☐ Cervical ☐ Vaginal ☐ Blood loss _____ cc			
Umbilical cord Clamping ☐ ☐ < 45s after delivery ☐ > 45s after delivery		Expression of umbilical cord ☐ Loose ☐ Tight ☐ Cut during delivery ☐ Cut after delivery	
Placenta ☐ Spontaneous ☐ Manual ☐ Umbilical vessels 2 3		Anomalies	
Complications None ☐ Meconium ☐ Fetal distress ☐ Fetal heart ☐ < 120 ☐ 120 - 160 ☐ Irregular ☐ Monitoring ☐ Clinic ☐ > 160 ☐ None ☐ Out-patient ☐ In-patient			
Signature of physician _____			Date _____

EVALUATION OF NEWBORN									
Sex ☐ ♂ ☐ ♀ ☐ Living ☐ Stillborn Mass _____ g		APGAR		0	1	2	1 min	5 min	10 Min
☐ Ophthalmic drops ☐ Vitamin K ☐ Baptized privately		Heart rate		Absent	Slow (below 100)	Over 100			
Nourishment ☐ Maternal ☐ Artificial ☐ homo. ☐ similia maternal		Resp. rate and effort		Absent	Slowing, irreg.	Good crying			
Resuscitation ☐ Yes ☐ No		Muscle tone		Flaccid	Flexion of ext.	Active motion			
Ventilation ☐ With sack and mask ☐ Tracheal tube _____ min ☐ Tracheal aspiration		Reflex irritability		No response	Grimace	Cry			
Asphyxia ☐ None ☐ Moderate ☐ Severe		Color		Blue - pale	Body pink ext. blue	All pink			
☐ Anomalies ☐ Complications		Remarks		Total					
Remarks _____ Parents informed ☐ Yes ☐ No		Name of physician caring for child _____		Date _____					

EVOLUTION OF THE MOTHER									
Post partum course ☐ Normal ☐ Abnormal Lowest HGB _____ Last HGB _____		Post partum haemorrhage ☐ Immediate ☐ Late Transfusion _____ sacks		Immunoglobuline anti-D Given _____ a m j par		Fever ☐ Yes ☐ No If yes, state as the case may be ☐ Endometritis ☐ Wound infection ☐ Other ☐ Other pelvic infection ☐ Thrombo-embolism ☐ Urinary infection ☐ Respiratory infection			
Condition on discharge Date yr m day _____ Alive ☐ Deceased ☐ Transferred ☐		Signature of physician _____		Date _____					

NOVA SCOTIA

FOR COPIES CONTACT : REPRODUCTIVE CARE PROGRAM, 5821 UNIVERSITY AVENUE, HALIFAX, NOVA SCOTIA B3H 1W3

[illegible]

DELIVERY RECORD

	TYPE OF LABOUR	INDUCTION	ANAESTHESIA
DATE OF BIRTH	SPONTANEOUS	FORCEPS	THIRD STAGE
HOUR OF BIRTH	WEIGHT	L.B./S.B./N.N.D.	EPISIOTOMY
PRES. POS.	BLOOD LOSS		NURSING
BABY-SEX	ASPHYXIA NEONATORUM		
COMPLICATIONS:			

DOCTOR'S SIGNATURE

POSTPARTUM EXAMINATION**DATE**

WEIGHT	LOCHIA		
B.P.	MENSES		
HAEMOGLOBIN	NURSING	PELVIS	PERINEUM
URINE	BREASTS	CERVIX	CORPUS
FAMILY PLANNING			

[illegible]

MATERNAL HISTORY & PHYSICAL




Physical Examination:		
Ht.:	Pres. Wt.	Wt. at LMP
Optic Fundi	Teeth	Thyroid
Heart	B.P.	ENT.
Lungs	Skin	
Breasts	Nipples	Tumors
Abdomen	Height of Fundus	
Fetal Heart	Presentation and Position	Amniotic Fluid Vol.
Extremities	Varicosities	Edema
Pelvic Exam:	CNS	Reflexes
Labour: ☐ Yes ☐ No ☐ Suspected		Risk Score
Summary Diagnoses, Problems and Management:		Low Risk = 0 - 2 High Risk = 3 - 6 Extreme Risk = $\geq$ 7
Date/Time		Signature

ANTE-NATAL RISK ASSESSMENT

I Reproductive History	II Associated Conditions	III Present Pregnancy
$< 16 = 1$ Age $16 - 35 = 0$ $> 35 = 2$ ☐		Bleeding < 20 Weeks $= 1$ ☐ ≥ 20 Weeks $= 3$ ☐
$0 = 1$ Parity $1 - 4 = 0$ $5+ = +2$ ☐	Previous Gynecological Surgery $= 1$ ☐ Chronic Renal Disease $= 2$ ☐ Gestational Diabetes $= 1$ ☐ Diabetes Mellitus $= 3$ ☐ Cardiac Disease $= 3$ ☐	Anemia < 10 GM % $= 1$ ☐ Prolonged Pregnancy (42 Weeks) $= 1$ ☐ Hypertension $= 2$ ☐ Premature Rupture Membranes $= 2$ ☐ Polyhydramnios $= 2$ ☐ Small For Dates $= 3$ ☐ Multiple Pregnancy or Breech or Malpresentation $= 3$ ☐ RH Isoimmunization $= 3$ ☐
Past OB History Habitual Abortion/Infertility $= 1$ ☐ P.P.H./Manual Removal $= 1$ ☐ Baby 9 lbs. (4082 gm.) $= 1$ ☐ Baby 5.5 lbs. (2500 gm.) $= 1$ ☐ P.E.T./Hypertension $= 1$ ☐ Previous Caesarean $= 2$ ☐ Stillbirth or Neonatal Death $= 3$ ☐ Prolonged Labour or Difficult Delivery $= 1$ ☐	Other Medical Disorders (Chronic Bronchitis, Lupus, etc.) ☐ Score according to Severity (1 to 3) ☐	
Category Score _____	Category Score _____	Category Score _____ Total Risk Score _____

Coopland & Al. Time

Low Risk 0 - 2
 High Risk 3 - 6
 Extreme Risk ≥ 7



/ RCP/01

ANTEPARTUM/INTRAPARTUM NURSING ADMISSION

Admission Date Time		Method of Admission ☐ Ambulatory ☐ Wheelchair ☐ Stretcher		Accompanied by _____ Is this your usual hospital of obstetrical care ☐ Yes ☐ No If no, referral by physician ☐ or self ☐																																																						
Reproductive History <table style="width: 100%;"> <tr> <td style="width: 15%;">Grav. </td> <td style="width: 15%;">LMP </td> <td style="width: 15%;">Prenatal Complications: ☐ Yes ☐ No Specify: _____</td> </tr> <tr> <td>Para </td> <td>EDC/dates </td> <td>Prev. Adm. this preg. ☐ Yes ☐ No When? _____</td> </tr> <tr> <td>Abort </td> <td>GEST. in wk. </td> <td>Prev. C/S ☐ Yes ☐ No Specify: _____</td> </tr> <tr> <td>S.B. </td> <td></td> <td></td> </tr> <tr> <td>N.D. </td> <td></td> <td></td> </tr> </table>						Grav.	LMP	Prenatal Complications: ☐ Yes ☐ No Specify: _____	Para	EDC/dates	Prev. Adm. this preg. ☐ Yes ☐ No When? _____	Abort	GEST. in wk.	Prev. C/S ☐ Yes ☐ No Specify: _____	S.B.			N.D.																																								
Grav.	LMP	Prenatal Complications: ☐ Yes ☐ No Specify: _____																																																								
Para	EDC/dates	Prev. Adm. this preg. ☐ Yes ☐ No When? _____																																																								
Abort	GEST. in wk.	Prev. C/S ☐ Yes ☐ No Specify: _____																																																								
S.B.																																																										
N.D.																																																										
Lab <table style="width: 100%;"> <tr> <td style="width: 20%;">Blood Grp. </td> <td style="width: 10%;">Date </td> <td style="width: 20%;">VDRL </td> </tr> <tr> <td>Rh Titre </td> <td> </td> <td>Hgb (Adm) </td> </tr> <tr> <td>Rh Serum </td> <td> </td> <td>Rubella Titre </td> </tr> </table> Date Where Pelvic X-Ray: ☐ Yes ☐ No Ultrasound: ☐ Yes ☐ No Other Tests: ☐ Yes ☐ No Specify: _____			Blood Grp.	Date	VDRL	Rh Titre		Hgb (Adm)	Rh Serum		Rubella Titre	Pertinent History <table style="width: 100%;"> <thead> <tr> <th></th> <th>Yes</th> <th>No</th> <th>Specify:</th> </tr> </thead> <tbody> <tr><td>Drug Allergies:</td><td>☐</td><td>☐</td><td></td></tr> <tr><td>Other Allergies:</td><td>☐</td><td>☐</td><td></td></tr> <tr><td>Medications:</td><td>☐</td><td>☐</td><td></td></tr> <tr><td>Non-Medical Drugs:</td><td>☐</td><td>☐</td><td></td></tr> <tr><td>Special Diet:</td><td>☐</td><td>☐</td><td></td></tr> <tr><td>Dentures:</td><td>☐</td><td>☐</td><td></td></tr> <tr><td>Contact Lenses:</td><td>☐</td><td>☐</td><td></td></tr> <tr><td>Glasses:</td><td>☐</td><td>☐</td><td></td></tr> <tr> <td>Smoker: ☐ Yes ☐ No</td> <td>Cig. Day in Preg. </td> <td>Cig. Day before Preg. </td> <td>No. of Yrs. </td> </tr> <tr> <td>Alcohol: ☐ Yes ☐ No</td> <td>Drinks Wk in Preg. </td> <td>Drinks Wk before Preg. </td> <td>No. of Yrs. </td> </tr> </tbody> </table>				Yes	No	Specify:	Drug Allergies:	☐	☐		Other Allergies:	☐	☐		Medications:	☐	☐		Non-Medical Drugs:	☐	☐		Special Diet:	☐	☐		Dentures:	☐	☐		Contact Lenses:	☐	☐		Glasses:	☐	☐		Smoker: ☐ Yes ☐ No	Cig. Day in Preg.	Cig. Day before Preg.	No. of Yrs.	Alcohol: ☐ Yes ☐ No	Drinks Wk in Preg.	Drinks Wk before Preg.	No. of Yrs.
Blood Grp.	Date	VDRL																																																								
Rh Titre		Hgb (Adm)																																																								
Rh Serum		Rubella Titre																																																								
	Yes	No	Specify:																																																							
Drug Allergies:	☐	☐																																																								
Other Allergies:	☐	☐																																																								
Medications:	☐	☐																																																								
Non-Medical Drugs:	☐	☐																																																								
Special Diet:	☐	☐																																																								
Dentures:	☐	☐																																																								
Contact Lenses:	☐	☐																																																								
Glasses:	☐	☐																																																								
Smoker: ☐ Yes ☐ No	Cig. Day in Preg.	Cig. Day before Preg.	No. of Yrs.																																																							
Alcohol: ☐ Yes ☐ No	Drinks Wk in Preg.	Drinks Wk before Preg.	No. of Yrs.																																																							
Reason for Admission				Labor ☐ Yes ☐ No (if yes) Prep ☐ Yes ☐ No Enema ☐ Yes ☐ No																																																						
Physical Assessment Pres. Wt. Wt. Gain Fundal Ht.		Abd. Palpation: Presentation Position Engagement		<table style="width: 100%;"> <tr> <td style="width: 10%;">T </td> <td style="width: 10%;">P </td> <td style="width: 10%;">BP </td> </tr> <tr> <td>R </td> <td>FH </td> <td>Reflexes </td> </tr> <tr> <td></td> <td></td> <td>Edema </td> </tr> </table>		T	P	BP	R	FH	Reflexes			Edema																																												
T	P	BP																																																								
R	FH	Reflexes																																																								
		Edema																																																								
Additional Information <table style="width: 100%;"> <tr> <td style="width: 20%;">Prenatal Classes:</td> <td>Yes ☐</td> <td>No ☐</td> </tr> <tr><td>Breast Feeding</td><td>☐</td><td>☐</td></tr> <tr><td>Rooming In:</td><td>☐</td><td>☐</td></tr> <tr><td>Keeping Baby:</td><td>☐</td><td>☐</td></tr> <tr><td>Seeing Baby:</td><td>☐</td><td>☐</td></tr> <tr><td>Social Worker Consult:</td><td>☐</td><td>☐</td></tr> </table>		Prenatal Classes:	Yes ☐	No ☐	Breast Feeding	☐	☐	Rooming In:	☐	☐	Keeping Baby:	☐	☐	Seeing Baby:	☐	☐	Social Worker Consult:	☐	☐	Nursing Plan/Comments High Risk Score		Patient's Requests: _____ Time Dr. Notified Dr. Signature _____																																				
Prenatal Classes:	Yes ☐	No ☐																																																								
Breast Feeding	☐	☐																																																								
Rooming In:	☐	☐																																																								
Keeping Baby:	☐	☐																																																								
Seeing Baby:	☐	☐																																																								
Social Worker Consult:	☐	☐																																																								

RCP/03

Time
hrs.☐ Intact

S.R.M.

RESEARCH

Quantity
of Fluid

Date	Date
------	------

Time

hrs.

hrs.[illegible]

Patient's Name and Hospital #		15	14	13	12	11	10	9	8	7	6	5	4	3	2	1	0
TIME	CONTENTS																
Medication																	
Drops/Min																	
Units/500 cc																	
Drugs IV Fluids Epidural																	
F.H.R.																	
B.P.																	
Pulse																	
Nurse's Signature																	

[illegible]

Transfer Time
to Postpartum _____

Condition _____

Signature _____

[illegible]

Fully Diluted: Date Time hrs.

Electronic Monitoring: Yes ☐ No ☐

2nd Stage if not monitored

F.H. Taken Last:	Rate	Time	hrs.
------------------	------	------	------

Summary of F.H.R. Tracing:

Nurse's Delivery Room Note:

Signature of Nurse

Mother's Companion: None ☐ Male ☐ Female ☐ Present During Labour ☐ Present At Delivery ☐

Neonatal Backup: Yes ☐ No ☐ Indication _____ Time Called _____ hrs. Time Arrived _____ hrs.

Catheterized: Yes ☐ No ☐ Time hrs.

Sutures: Yes ☐ No ☐ Type _____ sponge Count Correct Yes ☐ No ☐

Needle Count Correct Yes ☐ No ☐

Fill In If Applicable:

Doctor _____ Time Called _____ Time Arrived _____

Resident _____ Intern _____ Student _____

Anaesthetist _____ Resident Anaesthetist _____ Scrub Nurse _____

LABOR DELIVERY RECORD

LMP		Antepartum Status	ABO	Rh
E.D.C.		Grav.		BLOOD GR.
Gestn.		Para		Rh TITRE
At Del.		Abort		VDRL
☐ Singleton		S.B.		HGB
☐ Twins		N.N.D.		Rubella Titer
☐ Other				

Press Firmly

MEDICAL HISTORY

☐ Normal

☐ Abnormal Specify:

Problems Prev. Preg. or Infant

☐ No

☐ Yes Specify:

COMPLICATIONS OF THIS PREGNANCY

☐ Bleeding ☐ Toxemia ☐ None

☐ PROM ☐ Prem. Labour ☐ Diabetes

☐ Other Comment:

MEDICATIONS ☐ Yes ☐ No Specify:

SMOKING ☐ Yes ☐ No Cigs/Day

FIRST STAGE: Onset Date Time hrs. Duration

☐ Spontaneous ☐ Arm

☐ Induced ☐ Oxytocin Indication For

☐ Stimulated ☐ Prostaglandin

☐ Arm

☐ Oxytocin

☐ Prostaglandin

MEMBRANE RUPTURE Date Time

☐ Spont. ☐ Questionable ☐ Arm

☐ T. > 38 in Labour

☐ MECONIUM ☐ Yes ☐ No

Time First Noted

SECOND STAGE: Onset Date Time Duration

DELIVERY: Date Time

☐ Spontaneous ☐ Precipitate ☐ Operative

Position at Delivery

Forceps: ☐ Easy ☐ Moderate ☐ Difficult ☐ Trial only

☐ Outlet ☐ Low Mid ☐ Mid

FETAL HEART

☐ Normal ☐ Abnormal Specify:

Last Fetal pH Time

Last Maternal Blood Sugar Date Time

Rotation: ☐ Manual ☐ Forceps

Position at Interference

Position at Delivery

Breech: ☐ Spont. ☐ Assisted ☐ Forceps ☐ Extracted

Caesarean: ☐ Trans. ☐ Uterine Incision ☐ Vertical ☐ Hysterectomy

Episiotomy: ☐ None ☐ Midline ☐ Mediolateral

Lacerations: ☐ None ☐ Cervical ☐ Vaginal ☐ Anal Sphincter

Anal Mucosa ☐ Perineal ☐ Sutures

DRUGS TIME (Within 24 Hours) Drug/Dose/Route

THIRD STAGE: Completed Date Time hrs. Duration

Placenta Delivery Weight gm. Abnormalities:

☐ Spont. ☐ Assist. ☐ Manual

UMBILICAL VESSELS ☐ 3 ☐ 2

Oxytocin: ☐ None ☐ Before Separation ☐ After Separation

Type Dose Route

Blood Loss Before Del. After Del.

☐ Oxytocin Drip Postpartum P.P.H. ☐ Yes ☐ No

Anaesthesia:

☐ None ☐ Epidural ☐ Local ☐ General ☐ Pudendal ☐ Other ☐ Wedge used (15° Tilt)

Lowest B.P. Start Time

BABY: ☐ Girl ☐ Boy Weight gm.

☐ Liveborn ☐ Stillbirth: Date Last Fetal Movements Date Last Fetal Heart

CONDITION AT BIRTH: 1 min. 5 min.

Tone Colour Respiration Response Heart Rate

Age First Breath min. Age at First Cry min. Age Sust. Resp. min.

EYE MED'S

INFANT EXAM: ☐ Normal ☐ Abnormal Specify:

RESUSCITATION:

☐ None ☐ Oxygen Only ☐ Pos. Pres. Bag and Mask ☐ Pos. Pres. Bag and Tube ☐ Pos. Pres. Began Age to

Nurse Attending Delivered by Physician

MATERNAL POSTPARTUM FLOW SHEET

[illegible]

Tick (✓) or use symbol where appropriate
Use (X) where abnormal and describe on Progress Notes

RCP/05

**MATERNAL & NEWBORN
PROGRESS SHEET**[illegible]


**PHYSICIAN
NEWBORN ADMISSION & DISCHARGE**

Birth Date		Birth Weight (gm.)		Head Circ. (cm.)		Body Length (cm.)	
Clinical Gestational Assessment							
<36 wk.		36-37 wk.		38-40 wk.		41 = wk.	
Breast Tissue	☐ Absent	☐ 2mm	☐ 4mm	☐ 7mm			
Sole Creases	☐ Single Ant. Crease	☐ Anterior 1/3	☐ Few Posterior	☐ Full			
Ear Cartilage	☐ Nil	☐ Some	☐ Stiff				
Testes	☐ In Canal	☐ Descended					
Scrotum	☐ Few Rugae	☐ More					
				Gestational Age (Dates)		wk.	
				Gestational Assessment (Physical Exam)		wk.	
				Hypoglycemia Screening		Yes ☐ No ☐	
				Time			
				Dextrostix Results			
Admission Exam ☒ Normal ☒ Abnormal							
☐	General — Tone, Activity, Colour						
☐	Skin						
☐	Nutritional Status — Soft Tissue Wasting ☐ Yes ☐ No						
☐	Skull Shape						
☐	Fontanelle and Sutures						
☐	Eyes						
☐	Ears						
☐	Nose						
☐	Mouth						
☐	Palate						
☐	Respiratory System						
☐	Heart						
☐	Femoral Pulsations						
☐	Abdomen - Liver (.....cm.)						
	Spleen (.....cm.)						
☐	Umbilicus						
☐	Anus						
☐	Genitalia						
☐	Neurological						
☐	Skeletal, Clavicle Hips Feet						
Date		Signature					
Comments							
Discharge Exam ☒ Yes Comments if Abnormal							
☐	Tone and Activity						
☐	Skin — Jaundiced ☐ Yes ☐ No						
☐	Heart						
☐	Respiratory						
☐	Umbilicus						
☐	Circumcision						
☐	Feeding ☐ Breast ☐ Bottle Formula Type						
☐	Discharge Weight gm.						
☐	Blood Group						
☐	Hgb						
Discharge Diagnosis							
☐	Normal						
☐	Other						
Date		Signature					
Comments							
Date		Signature					

NEWBORN T.P.R.

Date															
Newborn															
Day	Q 1h x 4				Q 4h x 3				Q 8h						
Hour															
Temperature •	40													40	
	39													39	
	38													38	
	37													37	
	36													36	
	35													35	
	Pulse x	200													200
		190													190
		180													180
		170													170
160														160	
150														150	
140														140	
130														130	
120														120	
110														110	
Respirations •	100													100	
	90													90	
	80													80	
	70													70	
	60													60	
	50													50	
	40													40	
	30													30	
	20													20	


NEWBORN WEIGHT GRAPH




NEWBORN NURSING NOTES

[illegible]

ATLANTIC NEWBORN MALE GROWTH CHART



ATLANTIC NEWBORN FEMALE GROWTH CHART



NEWFOUNDLAND AND LABRADOR PRE-NATAL RECORD

HOSPITAL

NAME

ADDRESS

RELIGION

TELEPHONE

D.O.B.

AGE

OCCUPATION

ETHNIC GROUP

HUSBAND'S NAME

AGE

OCCUPATION

DOCTOR'S NAME

TOPICS FOR DISCUSSION

PRE-NATAL
CLASSES

VITAMIN
SUPPL.
DIET AND
WEIGHT

MENSTRUAL HISTORY

L.M.P.

E.D.C.

CYCLE

DURATION

NORMAL
SURE

ABNORMAL
UNSURE

BIRTH CONTROL

TYPE

NONE ☐

DATE DISCONTINUED

PREGNANCY TEST POSITIVE

(DATE)

PREVIOUS PREGNANCIES INCLUDING ABORTIONS

SMOKING

ALCOHOL

MEDICATIONS

LETTERS

BOOKS

EXERCISE

REST

COITUS

BOWELS

BIRTHS

GENETIC
COUNSELLING

DATE OF
ZUCKENING

BREAST FEED

DATE	PLACE	GEST.	WRS. OF LAB	TYPE OF DEL.	COMPLICATIONS	B.W.	SEX	CONDITION OF BABY & PRESENT HEALTH

PAST ILLNESSES

RENAL

☐

CARDIAC

☐

DIABETES

☐

INFECTIONS

☐

RUBELLA

☐

W.D.

☐

OPERATIONS

☐

TRANSFUSIONS

☐

EPILEPSY

☐

OTHER

☐

CURRENT MEDICATIONS

ALLERGIES

EXAMINATION

GENERAL CONDITION

WEIGHT

HEIGHT

HEAD AND NECK

CHEST

C.V.S.

BREASTS

ABDOMEN

EXTREMITIES (VARICOSITIES)

NEUROLOGICAL

PELVIC EXAMINATION

VULVA

VAGINA

CERVIX

UTERINE SIZE

ADNEXA

HISTORY OF PRESENT PREGNANCY

BREAST CARE

VOMITING

☐

BLEEDING (WHEN)

☐

POSSIBILITY
OF C.S.

SMOKING

☐

ALCOHOL

☐

DRUG USE

☐

HOSP. ADM.
PROCEDURES

NUTRITION

☐

X-RAY

☐

OTHER

☐

ANAE'S AND
ANALG.

LABOUR
STAGES

DIABETES

☐

CARDIAC

☐

HYPERTENSION

☐

BABY CARE

T.B.

☐

TWINS

☐

GENETIC

☐

ACCOMMODATION

MALFORMATIONS

☐

OTHER

☐

FAMILY HISTORY

WHITE COPY: MOTHER'S HOSPITAL CHART

YELLOW COPY: PHYSICIAN'S COPY

Please print firmly.

CLINICAL ASSESSMENT OF PELVIS	HOSPITAL INFORMATION		
	BREAST FEEDING	PRE-NATAL CLASSES	PREP TYPE
	ROOMING IN	HUSBAND IN	
SUMMARY OF DELIVERY	POST-PARTUM VISIT		DATE

A GUIDE TO PREGNANCY RISK GRADING

The risk factors or problems listed below are intended as examples only and the list is not exclusive. This risk grading system is intended as a basis for planning ongoing management of the pregnancy.

GRADE C Pregnancy at high risk

Pregnancies which are so complicated that the fetus and/or mother are obviously in danger. Wherever possible, these patients should be transferred to a regional hospital for intensive care and delivery.

DIABETES CLASS C, D, F, R, OR SIGNIFICANTLY COMPLICATED
RENAL DISEASE WITH HYPERTENSION
• 1 FUNCTION
PREMATURE RUPTURE OF MEMBRANES
(• SEPSIS)
PROLONGED RUPTURE OF MEMBRANES MORE THAN 6 HOURS, ANTEPARTUM BLEEDING, CONTINUING OR REPEATED*

HYPERTENSION WITH SUPERIMPOSED TOXAEMIA
EARLY UNCONTROLLED PREMATURE LABOUR*
SEVERE FETAL GROWTH ARREST
(• 10th Percentile)
HEART DISEASE, ESPECIALLY WITH FAILURE SEVERE PREECLAMPSIA.
* particularly 24 - 34 wks gestational age.

GRADE B Pregnancy at risk

The fetus and/or mother are definitely at risk and consultation should be obtained with a specialist obstetrician in your area. A plan of management will be suggested by him which may include collaborative care for the rest of the pregnancy with delivery usually in the regional hospital.

DIABETES CLASS A (GESTATIONAL) OR CLASS B
HYPERTENSION WITHOUT TOXAEMIA
APH, CEASED AND IN HOSPITAL
CERVICAL INCOMPETENCE
HYDRAMNIOS
POST-DATE PREGNANCY (42 weeks +)
HISTORY OF PRIOR STILL BIRTH OR NEONATAL DEATH
MATERNAL OBESITY
SIGNIFICANT TOBACCO, ALCOHOL, DRUG INTAKE
RHESUS IMMUNIZATION
RENAL DISEASE WITHOUT HYPERTENSION


MILD TOXAEMIA
CONTROLLED PREMATURE LABOUR
MULTIPLE PREGNANCY
BREECH PRESENTATION OR MALPRESENTATION
PRIMIGRAVIDA (age 15 and below, 35 and above)
HISTORY OF GENETIC OR METABOLIC DISEASE IN FAMILY (Genetic Amniocentesis or Counselling required)
ANAEMIA NOT RESPONDING TO IRON (• 10 gm)
WEIGHT GAIN • 10 LBS BY 30 WEEKS
PREVIOUS GROWTH RETARDATION, PREMATURE LABOUR, OR CAESAREAN SECTION

GRADE A Pregnancy at no predictable risk

NO PRIOR PERINATAL MORTALITY OR LOW BIRTHWEIGHT INFANT
NO SIGNIFICANT MEDICAL DISEASE
NO PREGNANCY COMPLICATIONS NOW OR IN THE PAST
(Bleeding, Hypertension, Premature Labour)
FETAL GROWTH SEEMS ADEQUATE

GRADE N Newborn at risk

Delivery in a high risk center should be considered if there is reason to suspect probable neonatal problems based on past or family history.
PREVIOUS SEVERE RDS, SEIZURE DISORDERS, HIRSCHSPRUNG'S DISEASE, CYSTIC FIBROSIS, CONGENITAL ANOMALIES



INSTRUCTIONS:

1. Each antenatal visit, measure the symphysis-fundus height (measured with a tape and refer to graph on the left)
2. At each antenatal visit insert your current assessment of pregnancy risk grading (A, B, C or N) in the appropriate column
3. Two or more minor risk problems can combine to produce a major pregnancy risk
4. Use this grading as the basis for continuing management of the pregnancy, e.g.
C Consider transfer
B Consider consultation